

Spaces of Tilings, Finite Telescopic Approximations and Gap-Labelling

Jean Bellissard^{1,2}, Riccardo Benedetti³, Jean-Marc Gambaudo⁴

¹ Université Paul-Sabatier, 118 route de Narbonne, 31062 Toulouse, France

² Institut Universitaire de France

³ Dipartimento di Matematica, via F. Buonarroti 2, 56127 Pisa, Italy

⁴ Laboratoire de Topologie, U.M.R. 5584 du CNRS,
Université de Bourgogne, B.P. 47870- 21078 Dijon Cedex France

Abstract

For a large class of tilings of $\mathbb{R}^d$, including the Penrose tiling in dimension 2 as well as the icosahedral ones in dimension 3, the continuous hull Ω_T of such a tiling T inherits a minimal $\mathbb{R}^d$ -lamination structure with flat leaves and a transversal Γ_T which is a Cantor set. In this case, we show that the continuous hull can be seen as the projective limit of a suitable sequence of branched, oriented and flat compact d -manifolds. Truncated sequences furnish better and better finite approximations of the asymptotic dynamical system and the algebraic topological features related to this sequence reflect the dynamical properties of the $\mathbb{R}^d$ -action on the continuous hull. In particular the set of positive invariant measures of this action turns to be a convex cone, canonically associated with the orientation, in the projective limit of the d^{th} -homology groups of the branched manifolds. As an application of this construction we prove a *gap-labelling theorem*:

Consider the C^* -algebra $\mathcal{A}_T$ of Ω_T , and the group $K_0(\mathcal{A}_T)$, then for every finite $\mathbb{R}^d$ -invariant measure μ on Ω_T , the image of the group $K_0(\mathcal{A}_T)$ by the μ -trace satisfies:

$$\mathcal{T}_\mu(K_0(\mathcal{A}_T)) = \int_{\Gamma_T} d\mu^t \mathcal{C}(\Gamma_T, \mathbb{Z}),$$

where μ^t is the transverse invariant measure on Γ_T induced by μ and $\mathcal{C}(\Gamma_T, \mathbb{Z})$ is the set of continuous functions on Γ_T with integer values.

1 Introduction

It has been argued [4, 5] that the mathematical description of aperiodic solids can be made through the construction of the so-called *Noncommutative Brillouin Zone* (NCBZ) to replace the Bloch theory used for periodic crystals. In a recent work [8], the construction of this noncommutative manifold was performed from the equilibrium positions of the atoms of the solid, seen as a *uniformly discrete* subset of the physical space $\mathbb{R}^d$. A suitable compactification of the family of $\mathbb{R}^d$ -translates of this set leads to the notion of *Hull*, which is a compact space Ω endowed with an action of $\mathbb{R}^d$ by homeomorphisms. The C^* -algebra of continuous functions on the NCBZ is then given by the *crossed product* [28] $\mathcal{C}(\Omega) \rtimes \mathbb{R}^d$ of the algebra of the Hull by the $\mathbb{R}^d$ -action.

The Hull can also be seen as a *lamination* [16] or a *foliated space* [27], namely a foliation with non smooth transverse structure. On the other hand [34], the construction of the Voronoi cells from the point set of atomic positions, leads to a *tiling* of $\mathbb{R}^d$ by polyhedra touching face to face, from which the point set can be recovered by a dual construction. Hence, the construction of the Hull can equivalently be performed from three complementary point of views: as a dynamical system, as a lamination or foliated space, as a tiling. This latter point of view permits to select constraints using the tiling language more easily than using the language of point sets. The tilings that have mostly attracted the attention of experts are those with a finite number of patches with a bounded size modulo translations (the so called *finite pattern condition* or FPC) and satisfying *repetitivity*. This imposes severe constraints on the point set of atomic positions. In particular, repetitivity implies the minimality of the $\mathbb{R}^d$ -action, while the FPC implies that the lamination is transversally a Cantor set. It is fair to say however, that some tilings of interest have a finite number of tiles only if we identify tiles under both translation and rotation [32]: these are not considered in the present paper.

One of the main consequence of constructing the NCBZ is the so-called *Gap Labelling Theorem* [3, 4, 5]. If $\mathcal{A}$ is the C^* -algebra of continuous functions on the NCBZ, and if $H = H^*$ is a selfadjoint element of $\mathcal{A}$, any spectral gap of H can be associated with a canonical element of the K -group $K_0(\mathcal{A})$ [13]. Moreover, the $\mathbb{R}^d$ -invariant ergodic probability measures μ on the Hull are in one-to-one correspondence with the extremal *traces* on $\mathcal{A}$. Each such trace τ_μ canonically defines a group homomorphism $\tau_{\mu,*}$ from $K_0(\mathcal{A})$ into $\mathbb{R}$ the image of which is called the set of *gap labels*. Since $\mathcal{A}$ is separable it follows that

the set of gap labels is a countable subgroup of $\mathbb{R}$. It has been conjectured [8] that, if the Hull is completely disconnected transversally to the $\mathbb{R}^d$ action, then the set of gap labels is nothing but the $\mathbb{Z}$ -module generated by the integrals $\int_{\Omega} d\mu f$ whenever f runs through the set $\mathcal{C}(\Omega, \mathbb{Z})$ of integer valued continuous functions on the Hull. This conjecture was proved for the first time for $d = 1$ in [5] using the Pimsner-Voiculescu exact sequence [29]. The same method was used in [19] to extend the result for $d = 2$. In [7], the result for $d = 2$ was reestablished by using the Kasparov spectral sequence. In the case for which the Hull is given by a $\mathbb{Z}^d$ -action on a Cantor set X , Hunton and Forrest [17], using the technic of spectral sequences, have proved that the K -group is isomorphic to the group cohomology of $\mathbb{Z}^d$ with values in the group $\mathcal{C}(X, \mathbb{Z})$. While this result does not lead to the computation of the set of gap labels in general, it permits to compute the K -group in many practical situations that occur in physics [18, 20] as well as the set of gap labels for $d = 3$ [9]. The proof of this conjecture for tilings that satisfy repetitivity and FPC in an arbitrary dimension is one of the main results of this paper.

Remark 1.1 At the time this paper was written, Benameur and Oyono [10] on the one hand, Kaminker and Putnam [23] on the other, announced that they have proved the gap labelling conjecture for a $\mathbb{Z}^d$ -action on a Cantor set for any d . Our result goes beyond such situation.

In the present work, the construction of the Hull for tilings with a finite number of tiles modulo translations, is given in term of an inverse limit or *expanding flattening sequence* (EFS) of *branched oriented flat manifolds with dimension d* (BOF- d -manifolds), called the approximants. The existence of a flat metric on each BOF-manifold leads to the notion of parallel transport by constant vector fields and to an $\mathbb{R}^d$ -action on the inverse limit. Consequently, the $\mathbb{R}^d$ -action on the Hull is topologically conjugate to the $\mathbb{R}^d$ -action on inverse limit of the approximants. This construction was inspired by previous works on dynamical systems which can be seen as tilings in dimension 1 [38, 31, 21].

One consequence of this construction is the identification of the space of positive invariant measures on the Hull with the positive cone of the inverse limit of the top-homology groups of the approximants. This cone is canonically defined thanks to the orientation of the approximants. Hence, the set of positive invariant measures can be interpreted from various equivalent

point of views: as the positive traces on $\mathcal{A}$, as the positive cone of the d -homology of the Hull, or, as will be seen later, as *Ruelle-Sullivan currents* on the lamination.

The C^* -algebra of the noncommutative Brillouin zone can be constructed from the inductive limit of the space of continuous functions on the approximants. Consequently, thanks to the Thom-Connes theorem, the K_0 -group of the NCBZ can be seen as the inductive limit of the K_0 -group of the approximants. For a BOF- d -manifold, the K -theory can be computed from a spectral sequence very similar to the one constructed for $\mathbb{Z}^d$ -action on a Cantor set [17]. Moreover, the set of gap labels can also be obtained from the so-called *Ruelle-Sullivan current* of the lamination, which in turn is shown in this paper to be given by a compatible family of d -cycles defined on each BOF-manifold of the EFS defining the lamination. The explicit construction of such family is given here and permits to prove the gap labelling theorem for the class of tilings under investigation.

1.1 Tilings

This subsection is devoted to few important properties of tilings in the usual d -dimensional Euclidean space $\mathbb{R}^d$. For any x in $\mathbb{R}^d$ and any $\epsilon > 0$, $B_\epsilon(x)$ denotes the closed d -ball in $\mathbb{R}^d$ with center x and radius ϵ and $\|\cdot\|$ the Euclidean norm in $\mathbb{R}^d$. The vector space $\mathbb{R}^d$ is *oriented* by stipulating that the standard basis is positive. A *prototile* is a compact closed connected subset of $\mathbb{R}^d$ which is the closure of its interior¹. A prototile determines, up to direct isometries of $\mathbb{R}^d$, a *tile i -type* and up to translations of $\mathbb{R}^d$ only, a *tile t -type*. Once the kind of tile type i or t is fixed, a prototile will often be identified with its tile type. Let X be a *countable* collection of prototiles of pairwise distinct types (of a given kind i or t). A tiling T with tile types in X , consists of a countable collection of subsets of $\mathbb{R}^d$, $\{t_1, t_2, \dots, t_n, \dots\}$, called *tiles*, such that:

- each tile belongs to a tile type in X ;
- the union of the tiles $\cup_{i \geq 1} t_i$ covers the whole space $\mathbb{R}^d$;
- the interiors of the tiles are pairwise disjoint.

¹It is often assumed in the literature that prototiles are homeomorphic to closed ball. The generalization we impose here is motivated by further constructions, but the standard results we will recall, work as well in our situation.

A tiling T is *polyhedral* if all its tiles are polyhedra in $\mathbb{R}^d$.

An example of polyhedral tiling is the Voronoi tiling of a Delone set [25]. A countable subset $\mathcal{L}$ of $\mathbb{R}^d$ is *uniformly discrete* if there is $r > 0$ such that every open ball of radius r meets $\mathcal{L}$ on at most one point. $\mathcal{L}$ is *relatively dense* if there is $R > 0$ such that every open ball of radius R meets $\mathcal{L}$ on at least one point. $\mathcal{L}$ is a *Delone set* if it is both uniformly discrete and relatively dense. If $x \in \mathcal{L}$, the *Voronoi cell* V_x is the open set of points in $\mathbb{R}^d$ that are closer to x than any other points in $\mathcal{L}$. If $\mathcal{L}$ is a Delone set, each Voronoi cell contains an open ball of radius r and is contained into an open ball of radius R . Moreover, the family $\{\bar{V}_x; x \in \mathcal{L}\}$ defines a polyhedral tiling of $\mathbb{R}^d$, in which the tiles are convex and meet face-to-face. Conversely, let X be a set of prototiles with pairwise distinct tile types, such that each prototile contains a ball of radius r and is contained in a ball of radius R . Define in each prototile p of X a point x_p , namely X is *punctured*. Then if T is a tiling with type tiles in X , the family of points x_t associated with each tile t by identification of t with a prototile, is a Delone set. However, the corresponding Voronoi tiling does not usually coincide with T .

Given a countable set X of prototiles, it is not known in general whether the set of tilings with type tiles in X is empty or not. However, many examples of finite X 's are known to give a rich set of tilings. For example, tiles of the Penrose tiling or the Voronoi tiling of a quasi crystal [34], belong to finitely many t -types in X , while tiles of the pinwheel tiling [32] belong only to two tile i -types but to an infinite number of tile t -types.

In this paper we focus our attention on the following type of tiling spaces:

- *we work with tile t -type and assume that the set of prototiles X is finite;*
- *we consider the tiling space $T(X)$ made with tilings for which the tile t -type of each tile is in X ;*
- *furthermore, it will be always assumed that $T(X) \neq \emptyset$.*

The group $(\mathbb{R}^d, +)$ acts on $T(X)$ by translations:

$$\omega : \mathbb{R}^d \times T(X) \rightarrow T(X) \quad \omega(x, T) = T + x.$$

The tiling space $T(X)$ is endowed with a distance (hence with the induced topology) defined as follows: let A denote the set of $\epsilon \in]0, 1[$ such that there exist x and x' in $B_\epsilon(0)$ such that $(T + x) \cap B_{1/\epsilon}(0) = (T' + x') \cap B_{1/\epsilon}(0)$, then:

$$\delta(T, T') = \inf A \quad \text{if } A \neq \emptyset$$

$$\delta(T, T') = 1 \quad \text{if } A = \emptyset .$$

Hence the diameter of $T(X)$ is bounded by 1 and the $\mathbb{R}^d$ -action on $T(X)$ is continuous.

For each $T \in T(X)$, $o(T) = T + \mathbb{R}^d$ denotes its *orbit*. The distance δ restricts to any orbit $o(T)$ and the induced topology is finer than the one induced by the Euclidean distance in $\mathbb{R}^d$; this topology reflects the way two translated tilings look the same at a local level. The *continuous Hull* of T , Ω_T , is the closure of $o(T)$ in $T(X)$. It is invariant for the $\mathbb{R}^d$ -action on $T(X)$.

Definition 1.2 A *patch* of $T \in T(X)$ is a finite sub-collection of tiles of T . The tiling T satisfies the *finite pattern condition* FPC if for any $s > 0$, there are *up to translation*, only finitely many patches with diameter smaller than s .

We refer to [25, 8] to understand, what means the FPC from the point set point of view.

Remark 1.3 Actually, the FPC is automatically satisfied in the case of a polyhedral tiling of $\mathbb{R}^d$, in which the prototiles are in finite number and meet face-to-face. However in the general case it is easy to construct tilings with a finite number of prototiles which do not satisfy the FPC.

This last definition is motivated by the following standard result (see the body and the references of [24]).

Proposition 1.4 *If a tiling $T \in T(X)$ satisfies the FPC, then Ω_T is compact so it coincides with the metric completion of $(o(T), \delta|_{o(T)})$.*

Remark 1.5 If the FPC holds, the metric on $T(X)$ induces the weak* topology defined in [8]. In other cases, while the tiling distance might lead to non compact continuous Hull, the weak* topology always lead to a compact Hull.

Notice that if a tiling T satisfies the finite pattern condition, then all tilings in Ω_T satisfy also the finite pattern condition.

Definition 1.6 Let X_p be a punctured copy of X , that is each prototile $p \in X$ is given with a marked point x_p in its interior. Consequently each tile t in a tiling $T \in T(X)$ admits a distinguished point x_t . The set Σ_{T_p} of tilings $T' \in \Omega_T$ such that one of the x_t coincides with the origin $0 \in \mathbb{R}^d$ is called the *canonical transversal* associated with X .

The following result is also standard [24] and admits an equivalent for Delone sets of finite types [8].

Proposition 1.7 *If a tiling $T \in T(X)$ satisfies the FPC, then for every punctured X_p , the canonical transversal Σ_{T_p} is compact and completely disconnected.*

Definition 1.8 A tiling $T \in T(X)$ satisfies the *repetitivity* condition if for any patch in T there exists $s > 0$, such that for every x in $\mathbb{R}^d$, there exists a translate of this patch which is in T and in the ball $B_s(x)$.

Again this last definition is motivated by the following proposition:

Proposition 1.9 *If a tiling $T \in T(X)$ satisfies both finite pattern and repetitivity conditions then the orbit $o(T')$ of any $T' \in \Omega_T$ is dense in Ω_T . In other words the dynamical system (Ω_T, ω) is minimal.*

Definition 1.10 A tiling $T \in T(X)$ is *aperiodic* if there exists no $x \neq 0$ in $\mathbb{R}^d$ such that $T + x = T$; it is *strongly aperiodic* if all tilings in Ω_T are aperiodic.

We have the following classical result (see [24]):

Proposition 1.11 *If an aperiodic tiling $T \in T(X)$ satisfies both finite pattern and repetitivity conditions then it is strongly aperiodic. In this case, any canonical transversal Σ_{T_p} is a Cantor set.*

Definition 1.12 The tilings $T \in T(X)$ which are aperiodic and satisfy both finite pattern and repetitivity conditions are said *perfect* tilings.

As the main object is the dynamical system (Ω_T, ω) , this suggests the following equivalence relation on tilings of $\mathbb{R}^d$.

Definition 1.13 Two tilings of $\mathbb{R}^d$, T and T' , are *Ω -equivalent* if there exists a homeomorphism $\phi : \Omega_T \rightarrow \Omega_{T'}$ which conjugates the two $\mathbb{R}^d$ -actions.

1.2 Main results

Let us describe now the content of the paper.

- In section 2 some basic general notions about *laminations* are given and, for a perfect tiling T of $\mathbb{R}^d$, it is proved that Ω_T can be equipped with a natural lamination structure whose leaves correspond to the $\mathbb{R}^d$ -orbits and which is transversely a Cantor set. Those laminations arising from perfect tilings are called *tilable* and are characterized. As an application we prove the following result:

Theorem 1.14 *Any perfect tiling is Ω -equivalent to some perfect tiling whose prototiles are d -rectangles.*

- In section 3 the category of branched oriented and flat (BOF) d -manifolds is defined. Local models, morphisms and in particular *BOF-submersions*, homology and cohomology are defined and studied.
- A particular class of projective limits of BOF- d -manifolds is defined in section 4.1. These projective limits are defined through BOF-submersions from one BOF- d -manifold to the previous one satisfying a *flattening property* introduced by Williams in [39]. The sequences that form the projective limits in this class are called *expanding flattening sequences* (EFS). Here again the homological and cohomological properties of these projective limits are described.
- In section 4.2 the connection between expanding flattening sequences and tilings is analyzed and leads to the following result:

Theorem 1.15 *Let T be a perfect tiling. Then its continuous Hull, seen as a dynamical system (Ω_T, ω) , is topologically conjugate to the inverse limit of an expanding flattening sequence of branched oriented flat manifolds, endowed with the action of $\mathbb{R}^d$ by parallel transport under constant vector fields.*

On one hand, an EFS defines a compact $\mathbb{R}^d$ -invariant subset of $T(X)$, for some finite set X of d -rectangles prototiles. Further conditions on the EFS ensure that this inverse limit is just one (perfect) Ω_T . On the other hand, for every perfect tiling T , the dynamical system (Ω_T, ω) can be realized as the inverse limit of a suitable EFS. Furthermore, not only the topological space Ω_T can be obtained by this inverse limit procedure but also most of the relevant structures on Ω_T . Hence, the $\mathbb{R}^d$ -action, the topological K -theory

groups $K^i(\Omega_T)$ and so on, can be obtained as inverse (or possibly direct) limits of suitable “pre-structures” on each of the BOF- d -manifolds forming the EFS. The “finite telescopic approximations” of (Ω_T, ω) mentioned in the title are the *truncated* finite subsequences of the EFS, decorated with their truncated sequences of pre-structures. These truncated sequences can be encoded by means of finite combinatorial patterns and provide a better and better approximation of the asymptotic dynamical system (Ω_T, ω) as the length of the truncated sequences grows up. The inverse limit approach to perfect *substitution tilings* [24] can be rephrased in terms of EFS and regarded as a special case of the previous construction. For substitution tilings the BOF- d -manifolds of the EFS are all equal each to the other, up to an overall rescaling.

- A particular attention is paid to the ergodic theory of tiling spaces in section 5. In the general case of a lamination with d -dimensional leaves, it is known that a transverse invariant measure μ^t can be seen as a Ruelle-Sullivan current i.e. a linear form on the vector space of d -forms on the lamination which satisfies some extra properties (see section 5). In the particular case of a perfect tiling space (Ω_T, ω) the correspondence between the transverse invariant measures of the lamination, the invariant measures of the $\mathbb{R}^d$ -action and the invariant measure on a transversal under the action of the holonomy groupoid of the associated lamination are also recalled. Considering Ω_T as a projective limit of an EFS of BOF- d -manifolds B_i , the data of a finite $\mathbb{R}^d$ -invariant measure μ on (Ω_T, ω) correspond to a *positive weights system* μ_i on the d -cells of a natural cell decomposition of B_i , satisfying a Kirchoff-like law. An elementary, but important fact is that the orientation, which is part of the BOF structure, allows to interpret these weights systems also as a positive cone in the space of d -cycles on B_i . So it is possible to contract these cycles against any real d -cohomology class. This passes to the limit and the set of finite invariant measures of (Ω_T, ω) corresponds to the convex set of the projective limit of the positive cones of the d^{th} homology groups of the BOF- d -manifolds B_i . On the other hand, the d -cohomology group on Ω_T can be seen as the inductive limit of the d -cohomology groups of the BOF- d -manifolds B_i , each real d -cohomology class α on Ω_T is the direct limit of classes α_i on B_i . It follows that for every finite invariant measure μ , the following pairing can be defined:

$$\langle \mu | \alpha \rangle = \langle \mu_i | \alpha_i \rangle \in \mathbb{R},$$

for i big enough. The special case of integer valued cohomology classes is studied in more details in view of the gap-labelling theorem.

- Section 6 is devoted to the non-commutative aspects of tiling spaces theory with a special emphasis on a gap labelling theorem. The fundamental non-commutative geometric structure associated with the dynamical system (Ω_T, ω) is a suitable C^* -algebra $\mathcal{A}_T$. The topology of this non commutative manifold is given by the group $K(\mathcal{A}_T) = K_0(\mathcal{A}_T) \oplus K_1(\mathcal{A}_T)$ [13] (by Bott periodicity theorem $K_i \cong K_{i+2}$). The Thom-Connes theorem [13] shows that $K_i(\mathcal{A}_T)$ is isomorphic to the topological K -group $K^{i+d}(\Omega_T)$. On the other hand, with each finite $\mathbb{R}^d$ -invariant measure μ on Ω_T is associated a canonical *trace* on $\mathcal{A}_T$ which induces a group homomorphism $\mathcal{T}_\mu : K_0(\mathcal{A}_T) \cong K^d(\Omega_T) \rightarrow \mathbb{R}$. The study of the image of $\mathcal{T}_\mu$ is relevant in the framework of the so called gap labelling [3, 4, 5, 24, 8, 9]. The isomorphism $K_0(\mathcal{A}_T) \cong K^d(\Omega_T)$ allows to factorize the pairing between invariant measures and elements in $K_0(\mathcal{A}_T)$ has a pairing between a cycle (the invariant measure) and an integer valued cocycle (see section 6 for further precision), leading to:

Theorem 1.16 (Gap-labelling theorem) *Let (Ω_T, ω) be the dynamical system associated with a perfect tiling T . Let μ be a finite invariant measure, Γ_T be a transversal and μ^t the induced transverse invariant measure on Γ_T , then:*

$$\mathcal{T}_\mu(K_0(\mathcal{A}_T)) = \int_{\Gamma_T} d\mu^t \mathcal{C}(\Gamma_T, \mathbb{Z}),$$

where $\mathcal{C}(\Gamma_T, \mathbb{Z})$ is the set of continuous functions on Γ_T with integer values.

Remark 1.17 1)- The existence of a underlying lamination structure on Ω_T is implicit in the literature about tiling dynamics and, for instance, it is explicetely described in [16, 27]. The idea of getting finite approximations by means of suitable branched manifolds has, as a natural background, the encoding of measured laminations embedded into 2-dimensional (3-dimensional) manifolds by means of embedded “train-tracks” (“branched surfaces”) equipped with positive weight systems [39, 37]. In the realm of tiling dynamics, it has been already observed in [38] and used in [31, 21], that the dynamics of a minimal map on a Cantor set can be described as an inverse limit of oriented graphs. On the other hand, the dynamical systems

induced by one dimensional perfect tilings of $\mathbb{R}$ exactly corresponds to minimal dynamics on a Cantor set. Despite the fact that in dimension greater than 1, tilings are much more complicated objects, it turns out that this approach can be generalized by using the convenient category of branched oriented flat manifolds.

2)- In the present paper only tilings with a finite number of t -types are considered so that Ω_T is in a natural way a dynamical system with an action of $\mathbb{R}^d$ by translations. One could ask whether our constructions could be generalized to the case of tilings with a finite number of i -types and with hulls carrying an action of the full group of direct isometries of $\mathbb{R}^d$ could also be considered. We hope to face this problem in a forthcoming work.

Acknowledgments: It is a pleasure for J.-M. Gambaudo to thank I. Putnam for very helpful (e-mail) conversations, and in particular for having pointed out a mistake in a first version of the paper. He also thank the Department of Mathematics of the University of Pisa, where a part of this work has been done, for its warm hospitality. J. Bellissard is indebted to several colleagues, A. Connes, T. Fack, J. Hunton, J. Kellendonk, A. Legrand, who helped him becoming more familiar with the technicalities involved in the gap labelling theorem. He thanks M. Benameur and I. Putnam for letting him know their recent works [10, 23] prior to publication. He also wants to thank the MSRI (Berkeley) and Department of Mathematics of the University of California at Berkeley for providing him help while this work was done during the year 2000-2001.

2 Tilings versus Laminations

Let M be a compact metric space and assume there exist a cover of M by open sets U_i and homeomorphisms called *charts* $h_i : U_i \rightarrow V_i \times T_i$ where V_i is an open set in $\mathbb{R}^d$ and T_i is some topological space. These open sets and homeomorphisms define an atlas of a (d) -*lamination* structure with d -dimensional leaves on M , if the *transition maps* $h_{i,j} = h_j \circ h_i^{-1}$ read on their domains of definitions:

$$h_{i,j}(x, t) = (f_{i,j}(x, t), \gamma_{i,j}(t)),$$

where $f_{i,j}$ and $\gamma_{i,j}$ are continuous in the t variable and $f_{i,j}$ is smooth in the x variable. Two atlas are *equivalent* if their union is again an atlas. A

(d) -lamination is the data of a metric compact space M together with an equivalence class of atlas $\mathcal{L}$.

We call *slice* of a lamination a subset of the form $h_i^{-1}(V_i \times \{t\})$. Notice that from the very definition of a lamination, a slice associated with some chart h_i intersects at most one slice associated with another chart h_j . The *leaves* of the lamination are the smallest connected subsets that contain all the slices they intersect. Leaves of a lamination inherit a d -manifold structure, thus at any point in the lamination, it is possible to define the tangent space to the leaf passing through this point. A lamination is *orientable* if there exists in $\mathcal{L}$ an atlas made of charts $h_i : U_i \rightarrow V_i \times T_i$ and orientations associated with each V_i preserved by the restrictions $f_{i,j}$ of the transition maps to the leaves. $\mathcal{L}$ is *oriented* if one fixes one global orientation.

Definition 2.1 Given a lamination $(M, \mathcal{L})$, a *transversal* of $\mathcal{L}$ is a compact subset Γ of M such that, for any leaf L of $\mathcal{L}$, $\Gamma \cap L$ is non empty and is a discrete subset with respect to the d -manifold topology of the leaf L .

The laminations which are related to tilings are in fact of a very special kind.

Definition 2.2 An oriented (d) -lamination $(M, \mathcal{L})$ is *flat* if the following properties are satisfied:

1) there exists in $\mathcal{L}$ a maximal atlas made of charts $h_i : U_i \rightarrow V_i \times T_i$ such that the transition maps read:

$$h_{i,j}(x, t) = (f_{i,j}(x), \gamma_{i,j}(t)),$$

where the $f_{i,j}$ do not depend on the t variable and are restrictions to their domains of definitions of translations in $\mathbb{R}^d$.

Note that each leaf of $\mathcal{L}$ is equipped with a structure of oriented flat d -manifold inherited from the Euclidean structure on $\mathbb{R}^d$. Then the lamination satisfies the further condition:

2) Every leaf of $\mathcal{L}$, considered as an oriented flat d -manifold, is isometric to $\mathbb{R}^d$.

Definition 2.3 A lamination $(M, \mathcal{L})$ is said *tilable* if it is flat and possesses a transversal Γ which is a Cantor set.

Definition 2.4 We call a domain U of a chart $h : U \rightarrow V \times C$ of such a maximal atlas a *box* of the lamination. For any point x in a box B with coordinates $(p(x), c(x))$ in the chart h , the slice $h^{-1}(V \times c(x))$ is called the *horizontal* of x and the Cantor set $h^{-1}(\{p(x)\} \times C)$ is called the *vertical* of x in the box B .

Notice that these definitions make sense since the transition maps map a vertical on a vertical and a horizontal on a horizontal.

On a tilable lamination $(M, \mathcal{L})$ it is possible to define a parallel transport along each leaf of the lamination. We denote by $Par(M, \mathcal{L})$ the set of parallel vector-fields on $(M, \mathcal{L})$. This set is parameterized by $\mathbb{R}^d$. This gives a meaning to the notion of translation in the lamination by a vector u in $\mathbb{R}^d$, defining this way a dynamical system $(M, \omega_{\mathcal{L}})$. The lamination is *minimal* if such a dynamical system is minimal.

Consider a perfect tiling $T \in T(X)$ constructed from a finite collection of prototiles $X = \{p_1, \dots, p_k\}$. For each prototile p_i in X , fix one interior point y_i . The set $Y = \{y_1, \dots, y_n\}$ determines a punctured version X_Y of X . We denote by $\Omega_{T,Y}$ the canonical transversal associated to Y . From Proposition 1.11, we know that $\Omega_{T,Y}$ is a Cantor set. It is plain to check that the dependence on the set of marked points is continuous:

Proposition 2.5 *Let us denote by $F(\Omega_T)$ the set of compact subsets of Ω_T endowed with the Hausdorff distance. Then the map $Y \rightarrow \Omega_{T,Y}$ from $\prod_j \text{Int}(p_j)$ to $F(\Omega_T)$, is continuous.*

The topology of the Cantor set $\Omega_{T,Y}$ is generated by the countable collection of *clopen* sets $U_{T,P}$ where P is a patch of a tiling in $\Omega_{T,Y}$ which contains the origin 0 of $\mathbb{R}^d$ and $U_{T,P}$ consists in these tilings in $\Omega_{T,Y}$ whose restrictions to P coincide with P .

Theorem 2.6 *Let T be a perfect tiling. Then there is a minimal tilable lamination $(\Omega_T, \mathcal{L})$ such that:*

- 1) $(\Omega_T, \omega_{\mathcal{L}})$ and (Ω_T, ω) are conjugate dynamical systems.
- 2) *For any arbitrary choice of the finite set $Y = \cup_i \{y_i\}$ where each point y_i in the interior of each p_i 's, the set $\Omega_{T,Y}$ is a transversal of the lamination.*

Proof. Fix one $\Omega_{T,Y}$ and choose at tiling T' in $\Omega_{T,Y}$ and a patch P' of T' that contains in its interior, a tile containing 0. For each such pair (T', P')

we consider the clopen set $C(T', P')$ in $\Omega_{T,Y}$ which consists in the tilings of $\Omega_{T,Y}$ that coincide with T' on the interior of P' and we define

$$U(T', P') = T''' + v \quad \forall (v, T''') \in \text{int}(P') \times C(T', P').$$

We denote $\phi(T', P')$ the map:

$$\text{int}(P') \times C(T', P') \rightarrow U(T', P')$$

$$(v, T''') \mapsto T''' + v.$$

When T' runs over $\Omega_{T,Y}$ the $U(T', P')$'s form a cover of Ω_T from which we can extract a finite sub-cover, that we denote for sake of simplicity, $U_1, \dots, U_n$, denoting also $P_1, \dots, P_n$ the corresponding patches, $C_1, \dots, C_n$ the corresponding clopen sets and $\phi_1, \dots, \phi_n$ the corresponding maps. Each clopen set C_i can be decomposed in a collection of pairwise disjoint clopen sets $C_{i,j}$, $j = 1, \dots, k(i)$, with arbitrary small diameters. Since the tilings in Ω_T are perfect, we can choose these diameters small enough so that:

- For each $j = 1, \dots, k(i)$ and each $i = 1, \dots, n$,
the restriction of ϕ_i to $\text{int}(P_i) \times C_{i,j}$ is a homeomorphism onto its image that we denote by $U_{i,j}$;
- For each pair (i_1, i_2) in $\{1, \dots, n\}^2$ and for each j_1 in $\{1, \dots, k(i_1)\}$ there exists at most one j_2 in $\{1, \dots, k(i_2)\}$ such that $U_{i_1,j_1} \cap U_{i_2,j_2} \neq \emptyset$.
A tiling $\hat{T}$ in $U_{i_1,j_1} \cap U_{i_2,j_2}$ can be seen in two different ways: either as $T_1 + v_1 = \phi_{i_1}(v_1, T_1)$ or as $T_2 + v_2 = \phi_{i_2}(v_2, T_2)$ and it is important to notice that the vector $t_{1,2} = v_2 - v_1$ is independent of the choice of $\hat{T}$ in $U_{i_1,j_1} \cap U_{i_2,j_2}$. The domain of definition of the transition map $\phi_{i_2}^{-1} \circ \phi_{i_1}$ is $\text{int}(P_{i_1}) \cap (\text{int}(P_{i_2}) - t_{1,2}) \times C_{i_1,j_1} \cap (C_{i_2,j_2} - t_{1,2})$ and on this domain, the transition map $\phi_{i_2}^{-1} \circ \phi_{i_1}(v_1, T_1)$ reads:

$$\phi_{i_2}^{-1} \circ \phi_{i_1}(v_1, T_1) = (v_2, T_2) = (v_1 + t_{1,2}, T_1 + t_{1,2}).$$

It follows that Ω_T can be endowed with a structure of tilable lamination $\mathcal{L}$, where the leaf containing a tiling T' in Ω_T is the $\mathbb{R}^d$ -orbit $o(T')$ of T' . The set $\Omega_{T,Y}$ is a transversal of the lamination by construction and it is plain to check that any other $\Omega_{T,Y'}$ is also a transversal. A translation in

the lamination $(\Omega_T, \mathcal{L})$ defined using $Par(\Omega_T, \mathcal{L})$ coincides by construction with a usual translation acting on Ω_T , so the last statement is immediate and minimality is a direct consequence of the minimality of (Ω_T, ω) .

Remark 2.7 Recently L. Sadun and R.F. Williams [33] have proved that the continuous hull of a perfect tiling of $\mathbb{R}^d$ is homeomorphic to a bundle over the d -torus whose fiber is a Cantor set, a much nicer object than a lamination. Unfortunately, this homeomorphism is not a conjugacy: it does not commute with translations. However this shows that every tiling is orbit equivalent to a $\mathbb{Z}^d$ -action.

If at the first sight, tilings seem to be very rigid objects, their connections with laminations show that tiling spaces seen as dynamical systems are easier to handle with. This

possibility of working with laminations will in turn lead to new results on tilings. For this purpose and by analogy with Markov partitions in Dynamical Systems, let us introduce the notion of *box decomposition* of a tilable lamination.

Definition 2.8 A *well oriented d -rectangle* in $\mathbb{R}^d$ is a d -rectangle such that, for $0 \leq i \leq n$, each i -face is parallel to an i -space generated by i vectors of the canonical basis in $\mathbb{R}^d$. A *block* in $\mathbb{R}^d$ is a connected set which is a finite union of well oriented d -rectangles. A well oriented d -rectangle is a *well oriented d -cube* when it is a d -cube.

Definition 2.9 Consider a tilable lamination $(M, \mathcal{L})$.

Consider a box $B \subset M$ which reads $int(P) \times C$ in a chart h in $\mathcal{L}$ defined in a neighborhood of the closure $cl(B)$ where C is a clopen subset of a Cantor set.

- when P is a well oriented d -rectangle, B is called a *box of rectangular type*;
- when P is a block, B is called a *box of block type*.

Notice that these definitions are independent on the choice of the chart in $\mathcal{L}$.

Definition 2.10 For any box B (of block or rectangular type) which reads $(\text{int}(P) \times C)$ in a defining chart, the set $h^{-1}(\partial P \times C)$ is called the *vertical boundary* of the box B .

Lemma 2.1 Consider a finite collection boxes of block type $B_1, \dots, B_m$ of a tilable lamination. Then, there exists a finite collection of boxes of rectangular type in $B'_1 \dots B'_p$ such that:

- the B'_l 's are pairwise disjoint;
- the closure of $\cup_{l=1}^p B'_l$ coincides with the closure of $\cup_{j=1}^m B_j$;
- if a B'_l intersects a B_j then it is contained in this B_j .

Proof

Let us first prove this result with two boxes B_1 and B_2 . For $i = 1, 2$, let h_i be the chart defining B_i . We set $h_i(B_i) = \text{int}(P_i) \times C_i$. On its domain of definition, the transition map $h_{1,2}$ reads:

$$h_{1,2}(x, t) = h_1 \circ h_2^{-1} = (f_{1,2}(x), \gamma_{1,2}(t)),$$

where $f_{1,2}$ does not depend on the t variable and is the restriction of a translation in

$\mathbb{R}^d$ by a vector $t_{1,2}$. Since, for $i = 1, 2$, there exists a partition of the Cantor sets C_i in clopen sets $C_{i,j}$, $j = 1, \dots, k(i)$ with arbitrary small diameters, we can choose such a partition so that $\text{int}(P_1) \times C_{1,j_1} \cap h_{1,2}(\text{int}(P_2) \times C_{2,j_2})$ is the (possibly empty) product space $\text{int}(P_1) \cap (\text{int}(P_2) - t_{1,2}) \times \tilde{C}_1$ where $\tilde{C}_1$ is a clopen subset of C_{1,j_1} . Similarly $\text{int}(P_2) \times C_{2,j_2} \cap h_{2,1}(\text{int}(P_1) \times C_{1,j_1})$ is the (possibly empty) product space $\text{int}(P_2) \cap (\text{int}(P_1) + t_{1,2}) \times \tilde{C}_2$ where $\tilde{C}_2$ is a clopen subset of C_{2,j_2} . It follows that $h_1^{-1}(\text{int}(P_1) \times C_{1,j_1}) \cup h_2^{-1}(\text{int}(P_2) \times C_{1,j_2})$ is the reunion of five disjoint boxes of block type $A_1 \cup A_2 \cup A_3 \cup A_4 \cup A_5$, where

- $A_1 = h_1^{-1}(\text{int}(P_1) \times C_{1,j_1} \setminus \tilde{C}_1)$;
- $A_2 = h_1^{-1}(\text{int}(P_1) \setminus (\text{int}(P_2) - t_{1,2}) \times \tilde{C}_1)$;
- $A_3 = h_1^{-1}(\text{int}(P_1) \cap (\text{int}(P_2) - t_{1,2}) \times \tilde{C}_1)$;
- $A_4 = h_2^{-1}(\text{int}(P_2) \setminus (\text{int}(P_1) + t_{1,2}) \times \tilde{C}_2)$;

- $A_5 = h_2^{-1}(\text{int}(P_2) \times C_{2,j_2} \setminus \tilde{C}_2)$.

Since P_1 , P_2 , $P_1 \cap (P_2 - t_{1,2})$, $P_1 \setminus (P_2 - t_{1,2})$ and $P_2 \setminus (P_1 + t_{1,2})$ are blocks, they are also finite unions of well oriented d -rectangles $P'_1, \dots, P'_m$. This decomposition in d -rectangles induces a finite collection of disjoint rectangular boxes whose closure is $h_1^{-1}(\text{int}(P_1) \times C_{1,j_1}) \cup h_2^{-1}(\text{int}(P_2) \times C_{1,j_2})$. Applying the same procedure for all pairs (i_1, j_1) and (i_2, j_2) we get the desired finite collection of rectangular boxes. The result for a collection of n boxes in general is obtained by an easy induction.

Definition 2.11 A tilable lamination $(M, \mathcal{L})$ admits a *box decomposition of rectangular type* (resp. of *block type*) if there exists a finite collection of boxes of rectangular type (resp. of block type) $\mathcal{B} = \{B_1, \dots, B_n\}$ such that:

- the B_i 's are pairwise disjoint;
- the union of the closures of the B_i 's covers the whole lamination M .

Proposition 2.12 Any tilable lamination $(M, \mathcal{L})$ admits a *box decomposition of rectangular type*.

Proof. Let x be a point in M , U_i an open set in M containing x and $h_i : U_i \rightarrow V_i \times T_i$ a chart. Choose a small well oriented d -rectangle P_i in V_i and a small clopen set C_i in T_i such that $x \in U'_i = h_i^{-1}(\text{int}(P_i) \times C_i)$. Doing this for all x in M we construct a cover of M with the boxes of rectangular type U'_i . As M is compact, we can extract from this cover a finite one that we still denote $(U'_i)_{i=1, \dots, n}$. The proof is then a direct consequence of Lemma 2.1.

In terms of tilings, this last result implies that the tilings induced on each leaf of the lamination by the intersection of the leaf with the vertical boundaries of the boxes $B_1, \dots, B_n$ defined in Proposition 2.12 are made with at most n t -types of well oriented d -rectangles. In the case when the lamination is minimal, they are perfect. This is resumed in the next corollary which is a converse to theorem 2.6 :

Corollary 2.13 Consider a minimal tilable lamination $(M, \mathcal{L})$; then there exists a perfect tiling T of $\mathbb{R}^d$ made with a finite number of prototiles which are well oriented d -rectangles and such that $\Omega_T = M$ (in the sense of theorem 2.6).

As an immediate consequence of Theorem 2.6 and Corollary 2.13 we have (see definition 1.13):

Corollary 2.14 *For any perfect tiling T of $\mathbb{R}^d$ there exists a perfect tiling T' made with finite number of well oriented rectangular prototiles which is Ω -equivalent to T .*

This should recommend us to work only with perfect tiling T' made with a finite number prototiles which are well oriented d -rectangles. However we will perform construction on Ω -equivalent tilings that will force us to leave this category. For this reason from now on we shall focus on perfect tilings made with a finite number of prototiles which are blocks and, unless otherwise stated, we will work with box decomposition of block type .

Notice that there is a lot of freedom in the construction of the box decomposition of a tilable lamination done in Proposition 2.12.

Definition 2.15 Let $\mathcal{B}$ and $\mathcal{B}'$ be two box decompositions of block type of a same tilable lamination $(M, \mathcal{L})$. We say that $\mathcal{B}'$ is *zoomed out* of $\mathcal{B}$ if:

- (1) for each point x in a box B' in $\mathcal{B}'$ and in a box B in $\mathcal{B}$, the vertical of x in B' is contained in the vertical of x in B ;
- (2) the vertical boundaries of the boxes of $\mathcal{B}'$ are contained in the vertical boundaries of the boxes of $\mathcal{B}$;
- (3) for each box B' in $\mathcal{B}'$, there exists a box B in $\mathcal{B}$ such that $B \cap B' \neq \emptyset$ and the vertical boundary of B does not intersect the vertical boundary of B' .

Proposition 2.16 *Consider a minimal tilable lamination $(M, \mathcal{L})$. Then, for any box decomposition of block type $\mathcal{B}$, there exists another box decomposition of block type zoomed out of $\mathcal{B}$.*

Proof.

Step 0. Let $\mathcal{B} = \{B_1, \dots, B_n\}$ be a box decomposition of block type of M and for $i = 1, \dots, n$, let h_i be the chart mapping B_i on $P_i \times C_i$. As already observed in the proof of lemma 2.1, up to a splitting of the vertical of the point in each box, we can assume that the intersection $B_i \cap B_j$ for i, j in

$\{1, \dots, n\}$ (read in a chart) is the product (possibly empty) of a block by a clopen set.

Step 1. For each $i = 1, \dots, n$, we denote by $J(i)$ a set of l 's in $\{1, \dots, m\}$ such that $cl(B_l) \cap cl(B_i) \neq \emptyset$, and define:

$$W_i = \cup_{l \in J(i)} cl(B_l).$$

Let x be in B_i and consider in W_i the connected component of leaf of the lamination containing x , say L_x . For any subset K of $J(i)$ we denote by $B_{i,K}$ the box made of all these points x in B_i such that L_x intersects B_l if and only if l is in K . We get a partition:

$$B_i = \cup_{K \subset J(i)} B_{i,K}.$$

For each K in $J(i)$, we extend the box $B_{i,K}$ as follows:

$$B_{i,K}^{(1)} = \cup_{x \in B_{i,K}} int(L_x).$$

If the vertical sizes of the boxes of the box decomposition $\mathcal{B}$ are chosen small enough and since the leaves of the lamination are copies of $\mathbb{R}^d$, the open sets $B_{i,K}^{(1)}$ are boxes and for each $i = 1, \dots, n$, two boxes $B_{i,K}^{(1)}$ and $B_{i,K'}^{(1)}$ are disjoint when K and K' are disjoint subsets of $J(i)$. Consider the collection of boxes $B_{1,K}^{(1)}$ when K runs over the subsets of $J(1)$ and, to get simpler notations, call this sequence $B_1^{(1)}, \dots, B_m^{(1)}$.

For $l = 2, \dots, n$, consider the boxes $B_{m+l-1}^{(1)} = B_l \setminus (\cup_{k=1}^{m+1} cl(B_k^{(1)}))$. It is clear that the sequence of boxes $\mathcal{B}^{(1)} = \{B_1^{(1)}, \dots, B_{m+n-1}^{(1)}\}$ is a box decomposition of the lamination M . This box decomposition satisfies points (1) and (2) of Definition 2.16. It is not (a priori) zoomed out of $\mathcal{B}$ since point (3) is satisfied only by the boxes $B_1^{(1)}, \dots, B_m^{(1)}$ which cover B_1 .

Step 2. We Consider now the box decomposition $\mathcal{B}^{(1)} = \{B_1^{(1)}, \dots, B_{m+n-1}^{(1)}\}$ and perform to the box $B_{m+1}^{(1)}$ the same procedure as the one we did in Step 1 for B_1 . We get this way a third box decomposition, $\mathcal{B}^{(2)} = \{B_1^{(2)}, \dots, B_{n_2}^{(2)}\}$ of the lamination M . This box decomposition satisfies points (1) and (2) of Definition 2.16 and point (3) is satisfied only by the boxes $B_1^{(1)}, \dots, B_m^{(1)}$ which cover B_1 and B_2 .

Step 3. We can iterate this procedure to get after n steps, a box decomposition $\mathcal{B}^n = \{B_1^{(n)}, \dots, B_{m_n}^{(n)}\}$ zoomed out of $\mathcal{B}$.

Consider two box decompositions $\mathcal{B} = \{B_1, \dots, B_n\}$ and $\mathcal{B}' = \{B'_1, \dots, B'_m\}$ of a same tilable lamination $(M, \mathcal{L})$ such that $\mathcal{B}'$ is zoomed out of $\mathcal{B}$. For each j in $\{1, \dots, m\}$, we denote by $I(j)$ the sets of i 's in $\{1, \dots, m\}$ such that $cl(B_i) \cap cl(B_j) \neq \emptyset$ and $B_i \cap B'_j = \emptyset$ and we define:

$$Z_j = \cup_{i \in I(j)} cl(B_i).$$

Definition 2.17 We say that $\mathcal{B}'$ *forces its border*² if for each j in $\{1, \dots, m\}$ and for each x in B'_j , the connected component L_x in Z_j of the leaf of the lamination containing x intersects $B_i \setminus B'_j$ if and only if i is in $I(j)$.

Actually Proposition 2.16 can be improved in the following theorem:

Theorem 2.18 *Consider a minimal tilable lamination $(M, \mathcal{L})$. Then, for any box decomposition of block type $\mathcal{B}$, there exists another box decomposition of block type zoomed out of $\mathcal{B}$ that forces its border.*

Proof. Fix j in $\{1, \dots, m\}$, and for each subset H of $I(j)$ we denote by $B'_{j,H}$ the box made of all these points x in B'_j such that L_x intersects $B_l \setminus B'_j$ if and only if l is in H . We get a partition:

$$B'_j = \cup_{H \subset I(j)} B'_{j,H}.$$

It is clear that the box decomposition made of all the $B'_{j,H}$'s, $H \subset I(j)$ and j in $\{1, \dots, m\}$ is a box decomposition which is zoomed out of $\mathcal{B}$ and forces its border.

Corollary 2.19 *Consider a minimal tilable lamination $(M, \mathcal{L})$. Then, for any box decomposition of block type $\mathcal{B}$, there exists a sequence of box decompositions of block type $\mathcal{B}_{n \geq 0}^{(n)}$ such that:*

- $\mathcal{B}^{(0)} = \mathcal{B}$;
- for each $n \geq 0$, $\mathcal{B}^{(n+1)}$ is zoomed out of $\mathcal{B}^{(n)}$ and forces its border.

²Actually the concept of "forcing its border" was first introduced in the context of tilings (see [24]). We will see in the sequel how our definition fits with the standard one.

In terms of tilings, this last result can be interpreted as follows. Consider a leaf L of the lamination, fix $n \geq 0$, and consider the tiling $T^{(n)}$ on this leaf induced by the intersection of the leaf with the vertical boundaries of the boxes of the box decomposition $\mathcal{B}^{(n)}$ defined in Theorem 2.18.

Definition 2.20 A sequence $T_{n \geq 0}^{(n)}$ of tilings obtained this way is called a *nested sequence of tilings*.

It satisfies in particular the following properties:
For all $n \geq 0$:

- the t -tiles types of $T^{(n)}$ coincide with a finite number of prototiles which are blocks;
- each tile $t^{(n+1)}$ of $T^{(n+1)}$ is a connected patch of $P^{(n)}$ of $T^{(n)}$ and contains at least one tile of $T^{(n)}$ its interior;
- furthermore, the patch $Q^{(n)}$ of $T^{(n)}$ made with all the tiles of $T^{(n)}$ that intersects $P^{(n)}$ is uniquely determined up to translation³;
- the tiling $T^{(n)}$ is Ω -conjugate to $T^{(0)}$.

Consider a minimal tilable lamination $(M, \mathcal{L})$ and a box decomposition $\mathcal{B}$. We define an equivalence relation on M as follows: two points in M are equivalent if they are in a same box of $\mathcal{B}$ and on the same vertical in this box. The quotient space is a branched d -manifold that inherits from its very construction some extra structures. Section 3 is devoted to an axiomatic approach of these objects. If we perform this quotient operation for a sequence of box decompositions zoomed out one of the other as in Corollary 2.19, we get a sequence of such branched manifolds whose study is reported in Section 4.

3 BOF-d-manifolds

In this section we will establish the basic notions concerning the category of *branched oriented flat* (briefly: BOF) d -manifolds. There exist several possible variations on these notions. We will try to adopt the simplest version suitable to applications to tilings.

³this is a generalization of the classical definition of "forcing its border" for substitution tilings (see [24]).

3.1 Local Models

For $r > 0$, and $x = (x_1, \dots, x_d)$ in $\mathbb{R}^d$, we denote by $\mathcal{C}(r, x)$ the d -dimensional open cube in $\mathbb{R}^d$ centered at x with size r :

$$\mathcal{C}(r, x) = \{(y_1, \dots, y_d) \mid \sup_{i=1, \dots, d} |y_i - x_i| < r\}.$$

For $i = 1, \dots, d$, let $\epsilon_i = \pm 1$ and consider the d -orthant $O_{\epsilon_1, \dots, \epsilon_d}(r, x)$ which consists in the closure of the points y in $\mathcal{C}(r, x)$ such that the sign of $y_i - x_i$ is ϵ_i for $i = 1, \dots, d$.

Definition 3.1 • A *type 1 d -cube* (with size $r > 0$, and centered at x) is the d -cube $(\mathcal{C}(r, x))$.

- For $1 \leq p \leq 2^d$, a *type p d -cube* (with size $r > 0$, and centered at x) is given by the data $(\mathcal{C}(r, x), (\mathcal{P}_1, n_1), \dots, (\mathcal{P}_p, n_p))$ where:

- (i) $\mathcal{P}_1, \dots, \mathcal{P}_p$ are p collections of d -orthants $\mathcal{P}_1, \dots, \mathcal{P}_p$ such that $\text{int}(\mathcal{P}_i) \cap \text{int}(\mathcal{P}_j) = \emptyset$ for $i \neq j$ and $\cup_{i=1}^p \mathcal{P}_i = \mathcal{C}(r)$;
- (ii) for $i = 1, \dots, p$, n_i is a positive integer associated to $\mathcal{P}_i$.

Consider the set of points $\mathcal{X} = \{(y, n)\}$ where y is in $\mathcal{C}(r, x)$ and $n = n_i$ whenever y is in $\mathcal{P}_i$. The corresponding type p d -cube (of size $r > 0$ centered at x) is the quotient space $\mathcal{X}/\sim$ where $(y, n) \sim (y', n')$ if and only if:

- $(y, n) = (y', n')$;
- or $y = y'$ and belongs to at least two distinct collections of d -orthants $\mathcal{P}_i$ and $\mathcal{P}_j$.

- A *BOF- d -cube* is a type p d -cube with some size $r > 0$ and centered at some x in RM^d , for some p in $1, \dots, 2^d$.

Consider a BOF- d -cube $\mathcal{D}$.

- The tangent plane $T_{\tilde{x}}\mathcal{D}$ to $\mathcal{D}$ at every point $\tilde{x}$ in $\mathcal{D}$ is well defined; thus the BOF- d -cube $\mathcal{D}$ has a natural *branched C^1 -structure*. In fact the *tangent bundle* $T(\mathcal{D}) = \cup_{\tilde{x}} T_{\tilde{x}}\mathcal{D}$ is trivial: one trivialization is associated with each positive basis of $\mathbb{R}^d$. We call *canonical trivialization* the trivialization associated with the standard basis of $\mathbb{R}^d$. In this way we have also specified one *orientation* of $\mathcal{D}$.

- The standard Euclidean metric on $\mathbb{R}^d$ ($ds^2 = dx_1^2 + \cdots + dx_d^2$) induces, in a natural way, a *branched flat metric* on $\mathcal{D}$, whence a distance, a notion of “branched geodesic arc” and so on.
- Each BOF- d -cube (with radius $r > 0$ and centered at x) is the union of a finite copies of a well oriented d -cube $\mathcal{C}(r, x)$ glued along collections of d -orthants, such that the orientation and both the branched C^1 -structure and the branched flat metric restrict to the usual ones. Any such an embedded well oriented d -cube $\mathcal{C}(r, x)$ is called a *smooth sheet* of the BOF- d -cube $\mathcal{D}$. A type p d -cube $\mathcal{D}$ is stratified as follows: for $1 \leq l \leq d$, $\mathcal{V}_l$ is the set of points (x, n) in $\mathcal{D}$ such that x belongs to exactly $d - l + 1$ distinct $\mathcal{P}_i$ ’s and $\mathcal{V}_0$ is the set of points (x, n) in $\mathcal{D}$ such that x belongs to strictly more than d distinct $\mathcal{P}_i$ ’s. We have:

- $\mathcal{D} = cl(\mathcal{V}_d) = \cup_{l=0}^{l=d} \mathcal{V}_l$;
- for each $0 < l \leq d$, $cl(\mathcal{V}_l) = \mathcal{V}_l \cup \mathcal{V}_{l-1}$ and $\mathcal{V}_l \cap \mathcal{V}_{l-1} = \emptyset$;
- $cl(\mathcal{V}_0) = \mathcal{V}_0$ is reduced to the center of $\mathcal{D}$.

For $1 \leq l \leq d$, each set $cl(\mathcal{V}_l)$ is a connected set which is a collection of l -faces of well oriented d -cubes containing the center of $\mathcal{D}$. Furthermore each point in $\mathcal{V}_l$ is the center of a type $d - l + 1$ d -cube in $\mathcal{D}$.

Definition 3.2 For $0 \leq l \leq d$ the set $\mathcal{V}_l$ is called the *l -stratum* of the BOF- d -disk $\mathcal{D}$.

.

Let us define now adapted morphisms between the local models.

Definition 3.3 A continuous map $f : \mathcal{D} \rightarrow \mathcal{D}'$ between two BOF- d -cubes of the same size $r > 0$ is a *local BOF-submersion* (onto its image) if:

- 1) f is C^1 with respect to the branched C^1 -structures.
- 2) For every open smooth sheet D of $\mathcal{D}$, $f(D)$ is a smooth sheet of $\mathcal{D}'$; the restriction of f to D coincides with the restriction of a translation. In fact, there is one translation which works for all smooth sheets of $\mathcal{D}$. If one fixes one basis of $\mathbb{R}^d$ and takes the corresponding trivializations of the two tangent bundles, then the differential df_x of f at any point x of $\mathcal{D}$, induces the identity on $\mathbb{R}^d$ (which has been identified with both $T_x \mathcal{D}$ and $T_{f(x)} \mathcal{D}'$).

Definition 3.4 A local BOF-submersion f is a *local BOF-isometry* iff it is bijective. In such a case $f(\mathcal{D})$ is a BOF- d -cube and f^{-1} is a local BOF-isometry and the stratification associated with $\mathcal{D}$ is mapped on the stratification associated with $f(\mathcal{D})$ (stratum by stratum).

3.2 BOF- d -manifolds

We are now in a position to extend our construction to global models.

Definition 3.5 A *BOF- d -manifold* B is a compact, connected metrizable topological space endowed with a (maximal) atlas $\{U_j, \phi_j\}$ such that:

- 1) every $\phi_j : U_j \rightarrow W_j$ is a homeomorphism onto an open set W_j of some open BOF- d -cube.
- 2) For any BOF- d -cube $\mathcal{D}$ embedded into $\phi_j(W_i \cap W_j)$, the restriction of $\phi_{ij} = \phi_i \circ \phi_j^{-1}$ to $\mathcal{D}$ is a local BOF-isometry onto a BOF- d -cube embedded into $\phi_i(W_i \cap W_j)$.

Hence for every point x of a BOF- d -manifold B , there exists a neighborhood U of x and a chart $\phi : U \rightarrow \phi(U)$ such that $\phi(U)$ is a BOF- d -cube and $\phi(x) = 0$. The type of the BOF- d -cube $\phi(U)$ is uniquely determined by x . Such a neighborhood U is called a *normal neighborhood* of x .

Definition 3.6 The *injectivity radius* of $x \in B$, denoted by $\text{inj}_B(x)$, is the sup of the size of any normal neighborhood of x in B . The *injectivity radius* of B , denoted $\text{inj}(B)$ is defined by:

$$\text{inj}(B) = \inf_{x \in B} \text{inj}_B(x) .$$

Definition 3.7 For $0 \leq l \leq d$, the *l -face* of a BOF- d -manifold B is the set of points x in B such that there exists a chart (U, ϕ) in the atlas such that x is in U and $\phi(x)$ belongs to the l -stratum of the BOF- d -cube $\phi(U)$. Notice that this property is independent on the choice of the chart (U, ϕ) as long as x is in U . A *l -region* is a connected component of the l -face of B . The finite partition of B in l -region, for $0 \leq l \leq d$, is called the *natural stratification* of B . The union of all the l -regions for $0 \leq l \leq d - 1$, forms the *singular locus* $\text{Sing}(B)$ of B .

Note that, in particular, any oriented flat d -torus B is a BOF- d -manifold with one d -region $R = B$. In general, any d -region of a BOF- d -manifold B is a connected (in general non compact) d -manifold naturally endowed with a $(X, G) = (\mathbb{R}^d, \mathbb{R}^d)$ -structure, where the group $G = \mathbb{R}^d$ acts on $X = \mathbb{R}^d$ by translation. This means that any region admits a d -manifold atlas such that all the transition maps on connected domains are restriction of translations. One can see for instance [11] for the basic notions about the (X, G) -structures set up.

Remark 3.8 All the objects that we have associated with any open BOF- d -cube (the tangent bundle, its trivializations, the orientation, the branched C^1 -structure, the branched flat metric and so on) globalize to any BOF- d -manifold. In particular, one has a natural notion of parallel transport on B with respect to the flat metric, whence the notion of *parallel vector field* on B . Let us denote by $Par(B)$ the set of parallel vector fields on B . Fix the canonical trivialization of the tangent bundle $T(B)$, then we have a natural isomorphism

$$\rho_B : \mathbb{R}^d \rightarrow Par(B)$$

defined by identifying $\mathbb{R}^d$ with any tangent space $T_{\tilde{x}}B$, and by associating to every vector $v \in \mathbb{R}^d$ the parallel vector field $\rho_B(v)$ obtained by parallel transport of v , starting from $\tilde{x}$. This does not depend on the choice of the base point $\tilde{x}$.

Definition 3.9 A BOF- d -manifold B has *rectangular faces* if, for each d -region R of B , there exists a C^1 injective map $f : R \rightarrow \mathbb{R}^d$ such that $f(R)$ is the interior of a well oriented d -rectangle in $\mathbb{R}^d$ and the differential of f , read in the charts of B , satisfies $df_x = Id$ at each point x in R .

In particular, the d -regions of a BOF- d -manifolds do not carry any topology.

Convention. From now on, we will only be concerned with BOF- d -manifolds with rectangular faces (unless otherwise stated).

3.3 BOF-submersion

Definition 3.10 A continuous map $f : B \rightarrow B'$ between BOF- d -manifolds is a *BOF-submersion* if:

- 1) f is C^1 and surjective.
- 2) For every $x \in B$ and for every normal neighborhood $\mathcal{D}'$ of $f(x)$ in B' , there exists a normal neighborhood $\mathcal{D}$ of x in B such that $f(\mathcal{D}) \subset \mathcal{D}'$ and, read in the corresponding charts, $f| : \mathcal{D} \rightarrow f(\mathcal{D})$ is a local BOF-submersion.
- 3) For each region R of B , there exists a region R' of B' such that f is a diffeomorphism from R' to R . In particular, the singular set of B is mapped into the singular set of B' :

$$\text{Sing}(B) \subset f^{-1}(\text{Sing}(B')).$$

Remark 3.11 Notice that the pre-image of a normal neighborhood U' in B' (with radius $r > 0$) is a finite union of disjoint normal neighborhoods $U_1, \dots, U_n$ in B (with radius $r > 0$) and that $f : U_i \rightarrow U'$ is a C^1 bijection whose differential is the identity when read in the charts. This imply in particular that:

$$\text{inj}(B') \leq \text{inj}(B).$$

3.4 Cycles and Positive Weight Systems

Definition 3.12 A (strictly) positive weight system w on a BOF- d -manifold B is a function which assigns to each d -region R of B a real number ($w(R) > 0$) $w(R) \geq 0$ in such a way that the “switching rules” (or Kirchoff-like laws) are satisfied; this means that along every $d - 1$ -region e of B the sum of the weights on the germs of d -regions along e on one side equal the sum of the weights of the germs of region on the other side. Let us denote by $(W^*(B))$ $W(B)$ the set of these weight systems. The BOF- d -manifold B equipped with $w \in W^*(B)$ is called a *measured* BOF- d -manifold. The *total mass* $m(w)$ of $w \in W(B)$ is just the sum of all the weights of w . So we have a partition $\{W^m(B)\}_{m \in \mathbb{R}^+}$ of $W(B)$ by the different total masses.

Remark 3.13 Note that the definition of positive a weight system does not involve any region orientation, and in fact it makes sense even for non-orientable branched manifolds (see for instance [12] for more details on this notion).

Let us now exploit the fact that the d -region orientations are part of the BOF- d -manifold structure. Let us first recall few elementary facts about the (cellular) (co)-homology of B . Consider the natural stratification of B , $\mathcal{V}_0, \dots, \mathcal{V}_d$, where, for $i = 0, \dots, d$, $\mathcal{V}_i$ is decomposed into a finite number a_i of i -region that we orient and order in an arbitrary way but for the orientation of the d -regions which is the natural orientation induced by the BOF d -manifold structure. We set $\mathbb{A} = \mathbb{Z}, \mathbb{R}$ and denote by $C_i(B, \mathbb{A})$ the free $\mathbb{A}$ -module which has as (ordered) basis the set of ordered and oriented i -regions in $\mathcal{V}_i$. By convention, for any oriented i -region e , $-e = -1e$, and it consists of the same cell with the opposite orientation. We call $C_i(B, \mathbb{A})$ the module of the i -chains of B . Clearly, as we dispose by definition of a distinguished basis of $C_i(B, \mathbb{A})$, it can be identified with $\mathbb{A}^{a_i}$.

Remark 3.14 With the exception of the d -regions orientation, there are no canonical choices for the above orders and orientations. On the other hand, it is clear that two different choices reflect in a very simple linear automorphism of $\mathbb{A}^{a_i}$ which is completely under control. Moreover, these choices will be essentially immaterial in our discussion.

We define the linear *boundary operator*

$$\partial_{i+1} : C_{i+1}(B, \mathbb{A}) \rightarrow C_i(B, \mathbb{A})$$

which assign to any $i+1$ -region, the sum of the i -regions that are in its closure pondered with a positive sign (resp. negative) if the induced orientation fits (resp. does not fit) with the orientation chosen for these i -regions. It is clear that $\partial_i \circ \partial_{i+1} = 0$.

The space $Z_i(B, \mathbb{A}) = \text{Ker } \partial_i$ is called the space of i -cycles of B and the space $B_i(B, \mathbb{A}) = \partial_{i+1}(C_{i+1}(B, \mathbb{A}))$ is called the space of i -boundaries of B . In fact, $B_i(B, \mathbb{A}) \subset Z_i(B, \mathbb{A})$ and $H_i(B, \mathbb{A}) = Z_i(B, \mathbb{A})/B_i(B, \mathbb{A})$ is the i^{th} homology group of B . Note that $H_d(B, \mathbb{A}) = Z_d(B, \mathbb{A})$.

A standard result of algebraic topology insures that (up to $\mathbb{A}$ -module isomorphism) $H_i(B, \mathbb{A})$ is a topological invariant of B that coincides with the i^{th} singular homology of B (see for example [35]). It is important to observe that a d -chain z of B is a d -cycle (i.e. $\partial_d(z) = 0$) if and only if the coefficients of z formally satisfy the above “switching rules” (or Kirchoff-like laws), extended to possibly non positive “weights”. Note that here we are using the distinguished choice of d -region orientations. Let us denote by $(Z_d^{>0}(B, \mathbb{A}))$ $Z_d^{\geq 0}(B, \mathbb{A})$ the (strict) positive cone of $Z_d(B, \mathbb{A})$, which is formed by the d -cycles with (strictly) positive coefficients. One clearly has

Lemma 3.15 $W(B) = Z_d^{\geq 0}(B, \mathbb{R})$, $W^*(B) = Z_d^{> 0}(B, \mathbb{R})$.

The dual module $C^i(B, \mathbb{A})$ of $C_i(B, \mathbb{A})$ is called the module of i -co-chain of B . It can be identified with $\mathbb{A}^{a_i}$ using the dual basis of the above distinguished basis of $C_i(B, \mathbb{A})$. There are linear *co-boundary* operators

$$\delta^i : C^i(B, \mathbb{A}) \rightarrow C^{i+1}(B, \mathbb{A})$$

defined by the relation $\delta^i(c)(z) = c(\partial_i(z))$. So we have the module of co-cycles $Z^i(B, \mathbb{A}) = \text{Ker} \delta^i$, the module of co-boundaries $B^i(B, \mathbb{A}) = \delta^{i-1}(C^{i-1}(B, \mathbb{A}))$, and finally (up to module isomorphism)

$$H^i(B, \mathbb{A}) = Z^i(B, \mathbb{A}) / B^i(B, \mathbb{A}) .$$

Note that $Z^d(B, \mathbb{A}) = C^d(B, \mathbb{A})$ which is the free $\mathbb{A}$ -module spanned by the *characteristic functions* of the d -regions of B .

If $\alpha \in H^d(B, \mathbb{A})$ is represented by a d -co-cycle $c = \sum_i s_i e_i^*$ (remind that we are using the dual basis), and $w = \sum_i w_i e_i$ is a d -chain, then

$$\alpha(w) = \langle w | c \rangle = \sum_i s_i \omega_i .$$

Remark 3.16 An immediate but relevant consequence is that one can *contract any d -co-homology class against a “positive measure”* on B (a fact which is not possible, for example, if the branched manifold is not orientable). For any $w \in W(B)$, it is not necessary to know that w is cycle in order to define $\langle w | c \rangle$, by using the last formula above. It should make sense even for a non-orientable branched manifold. What is remarkable in our situation is that $\langle w | c \rangle = \langle w | [c] \rangle$, where $[c]$ is the co-homology class represented by c . It follows that the natural pairing $\langle | \rangle$ induces a pairing

$$\langle | \rangle : W(B) \times H^d(B, \mathbb{A}) \rightarrow \mathbb{R} \quad \langle w | \alpha \rangle = \alpha(w) .$$

The following proposition is immediate.

Proposition 3.17 *If $f : B \rightarrow B'$ is a BOF-submersion then:*

1) there exists a natural linear map (well defined up to the mild ambiguity indicated in remark 3.14) $f_* : Z_d(B, \mathbb{A}) \rightarrow Z_d(B', \mathbb{A})$ such that $f_*(W^m(B)) \subset W^m(B')$, $f_*(W^{*,m}(B)) \subset W^{*,m}(B')$, for every $m \in \mathbb{R}^+$.

2) There exists a natural linear map $f^* : C^d(B', \mathbb{A}) \rightarrow C^d(B, \mathbb{A})$ such that $[\alpha](f_*(w)) = [f^*(\alpha)](w)$, for every $\alpha \in C^d(B', \mathbb{A})$, every $w \in Z_d(B, \mathbb{A})$, where $[\alpha] \in H^d(B, \mathbb{A})$ is the co-homology class represented by the co-chain α . In fact (with a slight abuse of notation), $f^*([\alpha]) = [f^*(\alpha)]$ well-defines a linear map $f^* : H^d(B', \mathbb{A}) \rightarrow H^d(B, \mathbb{A})$.

Lemma 3.18 *Let B be a BOF- d -manifold and $\mu = (\mu_1, \dots, \mu_k) \in W(B)$. Let $c \in Z^d(B, \mathbb{R})$ which represents a class of $H^d(B, \mathbb{Z})$. Then there exists a sequence of integers $(m_1, \dots, m_k)$ such that*

$$\langle \mu | c \rangle = \sum_i m_i \mu_i.$$

Moreover, all such linear combinations with integer coefficients arise in this way.

Proof. As μ is a d -cycle the value of $\langle \mu | c \rangle$ does not change if one add to c any co-boundary. As c represents a class in $H^d(B, \mathbb{Z})$, c differs from a suitable $c' \in Z^d(B, \mathbb{Z})$ by a co-boundary. The first statement of the lemma follows. The last one is a consequence of the fact that every d -co-chain in $C^d(B, \mathbb{Z})$ is actually a d -co-cycle.

4 Tilings & Expanding Flattening Sequences

4.1 Expanding Flattening Sequences

Definition 4.1 A BOF-submersion $f : B \rightarrow B'$ satisfies the *flattening condition*, if for every $x \in B$ and for every normal neighborhood $\mathcal{D}'$ of $f(x)$, there exists a small enough normal neighborhood $\mathcal{D}$ of x in B such that $f(\mathcal{D}) \subset \mathcal{D}'$ and, read in corresponding charts, $f| : \mathcal{D} \rightarrow f(\mathcal{D})$ is a local BOF-submersion that maps $\mathcal{D}$ on one single sheet of $f(\mathcal{D})$.

Definition 4.2 An *expanding flattening sequence* EFS is a sequence $\mathcal{F} = \{f_i : B_{i+1} \rightarrow B_i\}_{i \in \mathbb{N}}$ of submersions such that:

- (i) the sequence of injectivity radius of the B_i 's is a strictly increasing sequence that goes to $+\infty$ with i ;
- (ii) for each $i \in \mathbb{N}$ the map f_i satisfies the flattening condition.

Once an EFS $\mathcal{F}$ is chosen, comes, naturally associated with this sequence, the further “inverse” or “direct” sequences ($\mathbb{A} = \mathbb{Z}, \mathbb{R}$):

i)

$$Z_d(\mathcal{F}, \mathbb{A}) = \{(f_i)_* : Z_d(B_{i+1}, \mathbb{A}) \rightarrow Z_d(B_i, \mathbb{A})\}_{i \in \mathbb{N}}$$

with the restricted sequences

$$W(\mathcal{F}, \mathbb{A}) = \{(f_i)_* : W(B_{i+1}, \mathbb{A}) \rightarrow W(B_i, \mathbb{A})\}_{i \in \mathbb{N}}$$

and

$$W^*(\mathcal{F}, \mathbb{A}) = \{(f_i)_* : W^*(B_{i+1}, \mathbb{A}) \rightarrow W^*(B_i, \mathbb{A})\}_{i \in \mathbb{N}} .$$

ii)

$$C^d(\mathcal{F}, \mathbb{A}) = \{(f_i)^* : C^d(B_i, \mathbb{A}) \rightarrow C^d(B_{i+1}, \mathbb{A})\}_{i \in \mathbb{N}}$$

which induces

$$H^d(\mathcal{F}, \mathbb{A}) = \{(f_i)^* : H^d(B_i, \mathbb{A}) \rightarrow H^d(B_{i+1}, \mathbb{A})\}_{i \in \mathbb{N}} .$$

iii)

$$\text{Par}(\mathcal{F}) = \{df_i : \text{Par}(B_{i+1}) \rightarrow \text{Par}(B_i)\}_{i \in \mathbb{N}} .$$

Associated with these “inverse” (resp. “direct” sequences, come their *inverse limits* (resp. the direct limits) which will be relevant in the next sections and whose definitions are briefly recalled here.

Given an “inverse” sequence of maps $\mathcal{X} = \{X_i \xleftarrow{\tau_i} X_{i+1}\}_{i \in \mathbb{N}}$, let us recall that the elements of the *projective limit set* (or *inverse limit*)

$$\lim_{\leftarrow} \mathcal{X}$$

consists of the elements $(x_0, x_1, \dots, x_n, \dots)$ in the product $\prod_{i \geq 0} X_i$ such that

$$\tau_i(x_{i+1}) = x_i \text{ for all } i \geq 0.$$

Note that for every $j \geq 0$ there exists a natural map $p_j : \lim_{\leftarrow} \mathcal{X} \rightarrow X_j$.

If the X_j 's are topological spaces and the maps continuous maps, the set $\lim_{\leftarrow} \mathcal{X}$ is a topological space with the finest topology such that all the p_j 's are

continuous. Hence it is a subspace of the topological product $\prod_{i \geq 0} X_i$. Thus, if all the X_i 's are compact, $\lim_{\leftarrow} \mathcal{X}$ is compact.

Similarly, given any “direct” sequence of maps $\mathcal{Y} = \{Y_i \xrightarrow{\tau_i} Y_{i+1}\}$, the elements of the *direct limit set*

$$\lim_{\rightarrow} \mathcal{Y}$$

are the sequence of the form $(x_j, x_{j+1}, \dots, x_n, \dots)$, for some $j \geq 0$ such that $x_h \in Y_h$ and $\tau_h(x_h) = x_{h+1}$. Here again note that for every $j \geq 0$ there exists a natural map $i_j : Y_j \rightarrow \lim_{\rightarrow} \mathcal{Y}$. If the Y_j 's are topological spaces and the maps continuous, the direct limit set is a topological space with the roughest topology such that all the i_j 's are continuous.

Once these standard definitions have been recalled, let us consider again an EFS $\mathcal{F}$ together with the compact set $\Omega(\mathcal{F}) = \lim_{\leftarrow} \mathcal{F}$ and with the following associated inverse limits $\mathcal{M}^m(\mathcal{F}) = \lim_{\leftarrow} W^m(\mathcal{F}, \mathbb{R})$, $\mathcal{M}^{*m}(\mathcal{F}) = \lim_{\leftarrow} W^{*m}(\mathcal{F}, \mathbb{R})$. $\mathcal{M}^*(\mathcal{F}) = \cup_{m \in \mathbb{R}^+} \mathcal{M}^{*m}(\mathcal{F})$, $\mathcal{M}(\mathcal{F}) = \cup_{m \in \mathbb{R}^+} \mathcal{M}^m(\mathcal{F})$.

The next proposition is a consequence of standard facts on the co-homology of topological projective limits.

Proposition 4.3 $\lim_{\rightarrow} H^d(\mathcal{F}, \mathbb{A}) = H^d(\Omega(\mathcal{F}), \mathbb{A})$.

Since $\mathcal{M}(\mathcal{F}) = \lim_{\leftarrow} Z_d^{\geq 0}(\mathcal{F}, \mathbb{R})$, there is a natural pairing

$$\mathcal{M}(\mathcal{F}) \times H^d(\Omega(\mathcal{F}), \mathbb{R}) \rightarrow \mathbb{R} \quad < \mu | h > = h(\mu)$$

defined as follows :

For $h = (h_j, \dots, h_s, \dots)$ with $h_{s+1} = f_s^*(h_s)$, and $\mu = (\mu_0, \dots, \mu_s, \dots)$, then $< \mu | h > = < \mu_j | h_j >$.

The following result is a direct consequence of lemma 3.18.

Corollary 4.4 Let $\mu = (\mu_1, \dots, \mu_n, \dots)$ be in $\mathcal{M}(\mathcal{F})$, $c = (c_0, \dots, c_n, \dots)$ be a class in $H^d(\Omega(\mathcal{F}), \mathbb{Z})$ and let s be big enough so that $< \mu, c > = < \mu_s, c_s >$ then :

$$< \mu, c > = m_1 \mu_{s,1} + \dots + m_{p(s)} \mu_{s,p(s)},$$

where B_s has $p(s)$ d -regions, $\mu_{s,i}$ is the weights of the i^{th} d -region of B_s and the m_i 's are integers. Moreover, all such linear combinations with integer coefficients arise in this way.

In the next section we will show that $\Omega(\mathcal{F})$ is actually a space of tilings and that every continuous hull Ω_T can be obtained in this way. We will also see how $\omega_{\mathcal{F}} = \varprojlim \text{Par}(\mathcal{F})$ makes $\Omega(\mathcal{F})$ a dynamical system supporting an action of $\mathbb{R}^d$, which is conjugate to the usual action of $\mathbb{R}^d$ on tiling spaces. In section 5 we will identify $\mathcal{M}(\mathcal{F})$ with the set of invariant measures on $\Omega(\mathcal{F})$. In section 6 we will study the direct limit of the K -theory of the B_i . The application to the gap-labelling will follow.

4.2 From EFS to tilings

We associate with an EFS $\mathcal{F}$, the inverse limit $\omega_{\mathcal{F}} = \varprojlim \text{Par}(\mathcal{F})$.

Proposition 4.5 *The set $\omega_{\mathcal{F}}$ is naturally isomorphic to $\mathbb{R}^d$ and acts on $\Omega(\mathcal{F})$.*

Proof. Since, for any BOF- d -manifold B_i , there exists a natural isomorphism $\rho_{B_i} : \mathbb{R}^d \rightarrow \text{Par}(B_i)$ and for any BOF-submersion $f_i : B_{i+1} \rightarrow B_i$, the induced map $df : \text{Par}(B_i) \rightarrow \text{Par}(B_{i+1})$ is such that $(\rho_{B_{i+1}})^{-1} \circ df \circ \rho_{B_i} = \text{id}$, it follows that $\varprojlim \text{Par}$ is isomorphic to the inverse limit $\varprojlim \mathbb{R}^d$ associated with the inverse sequence $\text{Id} : \mathbb{R}^d \rightarrow \mathbb{R}^d$. Thus $\omega_{\mathcal{F}}$ is isomorphic to $\mathbb{R}^d$.

Let us show now that $\omega_{\mathcal{F}}$ acts on $\Omega(\mathcal{F})$. From Remark 3.11, we know that there exists $r > 0$ such that for each point $x = (x_0, \dots, x_i, \dots)$ in $\Omega(\mathcal{F})$ there exists a sequence of normal neighborhoods U_i with radius r around each point x_i in B_i such that for each $i > 0$, $f(U_i)$ is one single sheet D_{i-1} of U_{i-1} and thus $f(D_i) = D_{i-1}$. This gives a meaning to the notion of “small” translation of the point $(x_0, \dots, x_i, \dots)$. Finally any vector v in $\omega_{\mathcal{F}}$ can be

decomposed in a sum $v = \sum_{l=1}^{l=m} u_l$ where $\|u_l\| < r/2$ for $l = 1, \dots, m$. For a point x in $\Omega(\mathcal{F})$, we define $x + v = (\dots (x + u_1) + u_2) + \dots + u_m$. It is plain to check that this definition is independent of the decomposition.

Actually the dynamical system $(\Omega(\mathcal{F}), \omega_{\mathcal{F}})$ is a tiling space. Each region of the first BOF- d -manifold B_0 is a well oriented d -rectangle. Let X denote the set of prototiles made with all these well oriented d -rectangles. With any point $x = (x_0, \dots, x_n, \dots)$ in $\Omega(\mathcal{F})$ we associate a tiling $h(x)$ in $T(X)$ made with prototiles in X as follows:

- Since the injectivity radius of the B_i 's goes to infinity with i , there exists, for $i \geq 0$ a sequence of normal neighborhoods U_i of x_i with radius r_i where r_i is an increasing sequence going to infinity with i .
- From this sequence of normal neighborhoods we extract a sequence of (pre-images by chart maps of) sheets $D_i \subset U_i$ such that $D_i \subset f_i(D_{i+1})$. These sheets are well oriented d -cubes centered at x_i .
- Consider the translated copies $D'_i = D_i - x_i$. The d -cubes D'_i are centered at 0 in $\mathbb{R}^d$, they have an increasing radius going to infinity with i . Furthermore all these d -cubes are tiled with the prototiles in X and, for each $i > 0$, the tiling of D'_i coincides with the tiling of D'_{i-1} in D'_{i-1} .
- the limit of this process defines a single tiling $h(x)$ in $T(X)$.

The proofs of the following properties are plain.

Proposition 4.6 • *the map $h : \Omega(\mathcal{F}) \rightarrow T(X)$ is continuous, injective and conjugates the dynamical systems $(\Omega(\mathcal{F}), \omega_{\mathcal{F}})$ and $h(\Omega(\mathcal{F}), \omega)$ where ω stands for the restriction of the standard $\mathbb{R}^d$ -action on $T(X)$;*

• *by construction all tilings in $h(\Omega(\mathcal{F}))$ satisfy the finite pattern condition.*

As $\Omega(\mathcal{F})$ is compact, it is actually the union of continuous hulls of tiling orbits.

So far, we have not been concerned with the minimality of the dynamical system $(\Omega(\mathcal{F}), \omega_{\mathcal{F}})$; we give now a simple criterion that insures minimality. An EFS $\mathcal{F}$ verifies the *repetitivity condition* if for each $n \geq 0$ and each flat (pre-image by chart map of a) sheet D_i in B_i , there exists $p > 0$ such that each d -region of B_{i+p} covers D_i under the composition $f_i \circ f_{i+1} \circ \cdots \circ f_{i+p-1}$.

Proposition 4.7 *Let $\mathcal{F}$ be an EFS that satisfies the repetitivity condition then the dynamical system $(\Omega(\mathcal{F}), \omega_{\mathcal{F}})$ is minimal.*

Proof. The proof is exactly the same as the one for tilings (proposition 1.4).

So far we have shown how to associate with an EFS that satisfies the repetitivity condition, a compact space of perfect tilings made with rectangular prototiles, and the continuous hull of such a tiling. On the other hand, we have discussed in section 2, the correspondence between perfect tilings and minimal tilable lamination. In order to have a complete scheme of these correspondences it remains to us to be able to associate with any perfect tiling made with a finite number of well oriented rectangular prototiles, an EFS that satisfies the repetitivity condition.

4.3 From tilings to EFS

Consider a perfect tiling T made with the set of prototiles $X = \{p_1, \dots, p_n\}$ where the p_i 's are well oriented d -rectangles. We are going to associate with the hull Ω_T a first BOF- d -manifold and then an EFS that satisfies the repetitivity condition.

The First BOF- d -manifold B_0 is obtained by identifying two points in two different prototiles x_1 in p_1 and x_2 in p_2 if there exist a tile t_1 with t -type p_1 and a tile t_2 with t -type p_2 such that the translated copy of x_1 in t_1 coincides with the translated copy of x_2 in t_2 . There exists a natural projection $\pi_0 : \Omega_T \rightarrow B_0$ defined as follows: Consider a tiling T' in Ω_T , consider the prototile where the point 0 is and its location in the prototile. This defines a single point in B_0 . The projection π_0 also defines a box decomposition $\mathcal{B}^{(0)}$ (see Definition 2.9) of the hull equipped with its tilable lamination structure $(\Omega_T, \mathcal{L})$, where the sets $\pi_0^{-1}(int(p_i))$, for $i = 1, \dots, n$ are the n boxes of the box decomposition.

The BOF- d -manifold B_0 is a very poor approximation of the continuous hull Ω_T . In order to define better and better approximants of Ω_T , we construct a sequence of box decompositions $\mathcal{B}^{(i)}$, $i \geq 0$ such that for each $n \geq 0$, $\mathcal{B}^{(i+1)}$ is zoomed out of $\mathcal{B}^{(i)}$ and forces its border (see Corollary 2.19) and associate with this sequence a nested sequence of tilings $(T^{(i)})_{i \geq 0}$ where $T^{(0)} = T$ (see Definition 2.20). For $i \geq 1$, we construct the BOF- d -manifold B_i as follows:

- First we construct the BOF- d -manifold B'_i associated with $T^{(i)}$ in the same way we constructed B_0 associated with T_0 and a projection $\pi'_i : \Omega_T \rightarrow B'_i$.
- Since the tiles of $T^{(i)}$ are not well oriented d -rectangles but blocks and the BOF- d -manifold B'_i is not necessarily cellular.

- Each region of the BOF- d -manifold B'_i is tiled with the prototiles in X . The BOF- d -manifold B_i coincides B'_i but tiled with the prototiles in X . The d -regions of B_i are prototiles in X .

It is clear that for $i \geq 0$, B_i is a cellular BOF- d -manifold whose d -regions are well oriented d -rectangles, the map $\pi'_i : \Omega_T \rightarrow B'_i$ induces naturally a map $\pi_i : \Omega_T \rightarrow B_i$ and there exists a canonical map $f_i : B_{i+1} \rightarrow B_i$ that is defined by

$$f_i(x) = \pi_i(\pi_{i+1}^{-1}(x)).$$

Notice that this last definition makes sense since the vertical of a point of $(\Omega_T, \mathcal{L})$ in a box of $\mathcal{B}^{(i+1)}$ is included in its vertical in a box of $\mathcal{B}^{(i)}$ (see Definition 2.16). It is clear that, for each $i \geq 0$, the map $f_i : B_{i+1} \rightarrow B_i$ is a BOF-submersion and satisfies

$$f_i \circ \pi_{i+1} = \pi_i.$$

Furthermore, the sequence is an EFS since the flattening condition is a direct consequence of the fact that each box decomposition $\mathcal{B}^i$, for $i \geq 1$ forces its border (see for instance [24]). Let us denote by $\mathcal{F}$ the EFS we have gotten, $\Omega(\mathcal{F}) = \lim_{\leftarrow} \mathcal{F}$ and $\pi : \Omega_T \rightarrow \Omega(\mathcal{F})$ the map defined by $\pi(x) = (\pi_0(x), \pi_1(x), \dots, \pi_n(x), \dots)$.

Proposition 4.8 *The map $\pi : \Omega_T \rightarrow \Omega(\mathcal{F})$ is a conjugacy between (Ω_T, ω) and $(\Omega(\mathcal{F}), \omega_{\mathcal{F}})$.*

Proof. The proof is straightforward.

5 Invariant measures

So far we have described a same minimal dynamical system in 3 different ways:

- either as the continuous hull of a perfect tiling;
- or as a minimal tilable lamination;
- or as a minimal expanding flattening sequence.

The interplay between these 3 points of view is going to provide a combinatorial description of the invariant measures of this dynamical system. Before dealing with this very precise context, let us recall first some basics about transverse invariant measures for oriented laminations in the large (we refer to [16] for a more complete description).

Consider a lamination $(M, \mathcal{L})$ with d -dimensional leaves and fix an atlas in $\mathcal{L}$ made with the charts $h_i : U_i \rightarrow V_i \times T_i$ where V_i is an open set in $\mathbb{R}^d$ and T_i is some topological space. Recall that the transition maps $h_{i,j} = h_j \circ h_i^{-1}$ read on their domains of definitions:

$$h_{i,j}(x, t) = (f_{i,j}(x, t), \gamma_{i,j}(t)),$$

where $f_{i,j}$ and $\gamma_{i,j}$ are continuous in the t variable and $f_{i,j}$ is smooth in the x variable.

Definition 5.1 *Let $(M, \mathcal{L})$ be a lamination. A finite transverse invariant measure on $(M, \mathcal{L})$ is the data of a finite positive measure on each set T_i in such a way that if B is a Borelian set in some T_i which is contained in the definition set of the transition map γ_{ij} then:*

$$\mu_i(B) = \mu_j(\gamma_{ij}(B)).$$

It is clear that the data of a transverse invariant measure for a given atlas provides another invariant measure for any equivalent atlas and thus it makes sense to consider a transverse invariant measure μ^t of a lamination. The fact that the leaves of a lamination carry a structure of d -dimensional manifold allows us to consider differential forms on the lamination $(M, \mathcal{L})$. A k -differential form on $(M, \mathcal{L})$ is the data of k -differential forms on the open sets V_i that are mapped one onto the other by the differential of the transition maps f_{ij} . We denote by $A^k(M, \mathcal{L})$ the set of k -differential forms on $(M, \mathcal{L})$.

Definition 5.2 *A foliated cycle is a linear form from $A^d(M, \mathcal{L})$ to $\mathbb{R}$ which is positive on positive forms and vanishes on exact forms.*

There exists a simple way to associated with a transverse invariant measure a foliated cycle. Consider a d -differential form ω in $A^d(M, \mathcal{L})$ and assume for the time being, that the support of ω is included in one of the U_i 's. In this case, the form can be seen as a form on $V_i \times T_i$. By integrating ω on the

slices $V_i \times \{t\}$ we get a real valued map on T_i that we can integrate against the transverse measure μ_i to get a real number $\mathcal{C}_{\mu^t}(\omega)$. When the support of ω is not in one of the U_i 's, we choose a partition of the unity $\{\phi_i\}_i$ associated with the cover of M by the open sets U_i and define:

$$\mathcal{C}_{\mu^t}(\omega) = \sum_i \mathcal{C}_{\mu^t}(\phi_i \omega).$$

It is clear that we have defined this way a linear form $\mathcal{C}_{\mu^t} : A^d(M, \mathcal{L}) \rightarrow \mathbb{R}$ which does not depend on the choice of the atlas in $\mathcal{L}$ and of the partition of the unity. It is also easy to check that this linear form is positive for positive forms. The fact that $\mathcal{C}_{\mu^t}$ vanishes on closed form is a simple consequence of the invariance property of the transverse measure. The foliated cycle $\mathcal{C}_{\mu^t}$ is called *the Ruelle-Sullivan current* associated with the transverse invariant measure μ^t . It turns out that the existence of a foliated cycle implies the existence of a transverse invariant measure (see [36]) and thus both points of view: transverse invariant measure and foliated cycle are equivalent.

Let us consider now the more particular case of a tilable lamination $(M, \mathcal{L})$. Recall that on a tilable lamination it is possible to define a parallel transport along each leaf of the lamination which gives a sense to the notion of translation in the lamination by a vector u in $\mathbb{R}^d$, defining this way a dynamical system $(M, \omega_{\mathcal{L}})$. Let μ be a measure of finite mass on M invariant under the $\mathbb{R}^d$ -action. This invariant measure defines a transverse invariant measure of the lamination as follows. For any Borelian subset of a transverse set T_i :

$$\mu_i(B) = \lim_{r \rightarrow 0^+} \frac{1}{\lambda_d(V_i)} \mu(h_i^{-1}(V_i \times B)),$$

where λ_d stands for the Lebesgue measure in $\mathbb{R}^d$. Conversely, consider a transverse invariant measure μ^t of the tilable lamination $(M, \mathcal{L})$. Let $f : M \rightarrow \mathbb{R}$ be a continuous function and assume for the time being that the support of f is included in one of the U_i 's. In this case, the map $f \circ h_i^{-1}$ is defined on $V_i \times T_i$. By integrating $f \circ h_i^{-1}$ on the sheets $V_i \times \{t\}$ against the Lebesgue measure λ_d of $\mathbb{R}^d$, we get a real valued map on T_i that we can integrate again the transverse measure μ_i to get a real number $\int f d\mu$. When the support of f is not in one of the U_i 's, we choose a partition of the unity

$\{\phi_i\}_i$ associated with the cover of M by the open sets U_i and define:

$$\int f d\mu = \sum_i \int f \phi_i d\mu.$$

It is clear that we have defined this way a finite measure on M which does not depend on the choice of the atlas in $\mathcal{L}$ and of the partition of the unity and is invariant under the $\mathbb{R}^d$ -action. It is also plain that the existence of a finite measure on M invariant for the $\mathbb{R}^d$ -action is in correspondence with a finite measure on a transversal Γ invariant under the action of the holonomy groupoid.

Thus, for a tilable lamination $(M, \mathcal{L})$ the following 4 points of view are equivalent:

- A finite transverse invariant measure;
- a foliated cycle;
- a finite measure on M invariant for the $\mathbb{R}^d$ -action.
- a finite measure on a transversal Γ invariant for the holonomy groupoid action.

Let us now develop what follows for tilings from the above discussion. To fix the notations, we start with a perfect tiling T made with prototiles which are well oriented d -rectangles $X = \{p_1, \dots, p_q\}$. We associate with this tiling an EFS $\mathcal{F}$ constructed with a sequence of BOF-manifolds B_n , $n \geq 0$, and BOF-manifolds submersions: $f_n : B_{n+1} \rightarrow B_n$ that satisfy the flattening condition.

Let us denote by $\mathcal{M}(\Omega_T, \omega)$ the set of finite measures on Ω_T that are invariant under the $\mathbb{R}^d$ -action. We denote by $\mathcal{M}^m(\Omega_T, \omega)$ the set of finite measures on Ω_T with total mass m .

Choose a set Y made of one point in the interior of each prototiles. The transversal $\Omega_{T,Y}$ is a Cantor set on which acts the holonomy groupoid $\mathcal{H}_{T,Y}$. Let us denote by $\mathcal{M}(\Omega_{T,Y}, \mathcal{H}_{T,Y})$ the set of finite measures on $\Omega_{T,Y}$ that are invariant under the action of the holonomy groupoid $\mathcal{H}_{T,Y}$. With any finite invariant measure μ in $\mathcal{M}(\Omega(T), \omega)$ can be associated a finite *transverse* measure μ^t in $\mathcal{M}(\Omega_{T,Y}, \mathcal{H}_{T,Y})$ and this map is one to one. Since $\Omega_{T,Y}$ is a Cantor set, it can be cover by a partition in clopen sets with arbitrarily

small diameters. Such a partition $\mathcal{P}$ is *finer* than another partition $\mathcal{P}'$ if the defining clopen sets of the first one are included in clopen sets of the second one. Consider a sequence of partitions $\mathcal{P}_n$, $n \geq 0$ of $\Omega_{T,Y}$ such that, for all $n \geq 0$, $\mathcal{P}_{n+1}$ is finer than $\mathcal{P}_n$ and the diameter of the defining clopen sets of $\mathcal{P}_n$ goes to zero as n goes to $+\infty$.

Claim *A finite measure on $\Omega_{T,Y}$ is given by the countable data of non negative numbers associated with each defining clopen sets of each partition $\mathcal{P}_n$ which satisfy the obvious additivity relation.*

The EFS $\mathcal{F}$ provides us such a sequence of partitions $\mathcal{P}_n$ as follows. For each $n \geq 0$, let $F_1, \dots, F_{p(n)}$ be the regions of the BOF- d -manifold B_n . Each F_i is a copy of a prototile in X and consequently there is a marked point y_i in F_i . For $n \geq 0$, $i = 1, \dots, p(n)$, we consider the clopen set:

$$\mathcal{C}_{n,i} = \pi_n^{-1}(y_i).$$

For n fixed and when i varies from 1 to $n(p)$ the clopen sets $\mathcal{C}_{n,i}$ form a partition $\mathcal{P}_n$ of $\Omega_{T,Y}$. Furthermore, for $n \geq 0$, $\mathcal{P}_{n+1}$ is finer than $\mathcal{P}_n$ and the diameter of the clopen sets $\mathcal{C}_{n,i}$ goes to zero as n goes to $+\infty$. It follows that a finite measure on $\Omega_{T,Y}$ is given by the countable data of non negative weights associated with each defining clopen sets $\mathcal{C}_{n,i}$ which satisfy the obvious additivity relation.

The relation between an invariant measure μ in $\mathcal{M}(\Omega_T, \omega)$ and the associated transverse invariant measure μ^t in $\mathcal{M}(\Omega_{T,Y}, \mathcal{H}_{T,Y})$ is given by :

Proposition 5.3 *For $n \geq 0$ and i in $\{1, \dots, n(p)\}$:*

$$\mu^t(\mathcal{C}_{n,i}) = \frac{1}{\lambda_d(F_i)} \mu(\pi_n^{-1}(F_i)).$$

In the sequel, we are going to characterize in a combinatorial way these invariant measures.

Consider the map $\tau_n : \mathcal{M}(\Omega_T, \omega) \rightarrow C_d(B_n, \mathbb{R})$ defined by

$$\tau_n(\mu) = \left(\frac{\mu(\pi_n^{-1}(F_1))}{\lambda_d(F_1)}, \dots, \frac{\mu(\pi_n^{-1}(F_{p(n)}))}{\lambda_d(F_{p(n)})} \right),$$

where the regions $F_1, \dots, F_{p(n)}$ are now ordered and equipped with the natural orientation that allows to identify $C_d(B_n, \mathbb{R})$ with $\mathbb{R}^{p(n)}$.

Remark 5.4 Notice that the coordinates of $\tau_n(\mu)$ are the transverse measures associated with μ of clopen sets $\mathcal{C}_{n,io}$ for some $n \geq 0$ and some i in $\{1, \dots, n(p)\}$.

Proposition 5.5 For any $n \geq 0$, the map τ_n satisfies the following properties:

- (i) $\tau_n(\mathcal{M}(\Omega_T, \omega)) \subset W^*(B_n)$;
- (ii) $f_{n,*} \circ \tau_n = \tau_{n+1}$.

Proof.

(i) Choose an invariant measure μ in $\mathcal{M}(\Omega_T, \omega)$. The invariance of μ implies that on each edge of B_n the sum of the transverse measures associated with the regions on one side of the edge is equal to the sum of the transverse measures of regions on the other side of the edge. These are exactly the switching rules (or Kirchoff-like laws) that define $W(B_n)$. The fact that the measure is invariant implies that each region has a strictly positive weight. Thus $\tau_n(\mu)$ is in $W^*(B_n)$.

(ii) Let $F'_1, \dots, F'_{p(n+1)}$ be the ordered sequence of regions of B_{n+1} equipped with the natural orientation that allows to identify $C_d(B_{n+1}, \mathbb{R})$ with $\mathbb{R}^{p(n+1)}$. To the linear map $f_{n,*} : C_d(B_{n+1}, \mathbb{R}) \rightarrow C_d(B_n, \mathbb{R})$ corresponds a $n(p) \times n(p+1)$ matrix A_n with integer non negative coefficients. The coefficient $a_{i,j,n}$ of the i^{th} line and the j^{th} column is exactly the number of pre images in F'_j of a point in F_i . Thus we have the relations:

$$\frac{\mu(\pi_n^{-1}(F_i))}{\lambda_d(F_i)} = \sum_{j=1}^{j=p(n+1)} a_{i,j,n} \frac{\mu(\pi_{n+1}^{-1}(F'_j))}{\lambda_d(F'_j)},$$

for all $i = 1, \dots, p(n)$ and all $j = 1, \dots, p(n+1)$, which exactly means that $f_{n,*} \circ \tau_n = \tau_{n+1}$.

Let us set $\mathcal{M}^*(\mathcal{F}) = \varprojlim W^*$. Then we have the following characterization of the set of invariant measures of (Ω_T, ω) .

Theorem 5.6

$$\mathcal{M}(\Omega_T, \omega) \cong \mathcal{M}^*(\mathcal{F}).$$

Proof. The inclusion

$$\mathcal{M}(\Omega_T, \omega) \subset \mathcal{M}^*(\mathcal{F})$$

is a direct consequence of Proposition 5.5.

With any element $(\beta_0, \dots, \beta_n, \dots)$ in $\mathcal{M}^*(\mathcal{F})$ we can associate, thanks to Proposition 5.3 a weight to each clopen set $\mathcal{C}_{n,i}$. The relation $\beta_n = f_{\star n} \beta_{n+1}$ means that this countable sequence of weights satisfies the additivity property and then defines a measure on $\Omega_{T,Y}$. The fact all the β_n 's are cycles, i.e satisfy the switching rules, yields that this measure is a transverse invariant measure, i.e. an element in $\mathcal{M}(\Omega_{T,Y}, \mathcal{H}_{T,Y})$. Since the correspondence between $\mathcal{M}(\Omega_{T,Y}, \mathcal{H}_{T,Y})$ and $\mathcal{M}(\Omega_T, \omega)$ is bijective, the equality is proved.

Corollary 5.7 • *If the dimension of $H^d(B_n, \mathbb{R})$ is uniformly bounded by N , then for all $m > 0$, $\mathcal{M}^m(\Omega_T, \omega)$ contains at most N ergodic measures;*

- *if furthermore the coefficients of all the matrices $f_{\star n}$ are uniformly bounded then for all $m > 0$, $\mathcal{M}^m(\Omega_T, \omega)$ is reduced to a single point i.e; the dynamical system (Ω_T, ω) is uniquely ergodic.*

Proof. The proof is standard and can be found in [21] in a quite similar situation in the particular case when $d = 1$. To prove the first statement we may assume that the dimension of the $H^d(B_n, \mathbb{R})$'s is constant and equal to N . The set $\mathcal{M}^m(\Omega_T, \omega)$ is a convex set and its extremal points coincides with the set of ergodic measures. Since $\mathcal{M}^m(\Omega_T, \omega) = \mathcal{M}^{\star m}(\mathcal{F})$, the convex set $\mathcal{M}^m(\Omega_T, \omega)$ is the intersection of the convex nested sets:

$$\mathcal{M}(\Omega_T, \omega) = \bigcap_{n \geq 0} W_n$$

where

$$W_n = f_{\star 1} \circ \dots \circ f_{\star n-1} W^*(B_n).$$

Since each convex cone W_n possesses at most N extremal lines, the limit set $\mathcal{M}^m(\Omega_T, \omega)$ possesses also at most N extremal points and thus at most N ergodic measures.

In order to prove the second statement, we have to show that $\mathcal{M}(\Omega_T, \omega)$ is one dimensional. Consider two points x and y in the positive cone of $\mathbb{R}^N$. Let T be the largest line segment containing x and y and contained in the

positive cone of $\mathbb{R}^N$. We recall that the hyperbolic distance between x and y is given by:

$$Hyp(x, y) = -\ln \frac{(m+l)(m+r)}{l.r},$$

where m is the length of the line segment $[x, y]$ and l and r are the length of the connected components of $T \setminus [x, y]$. Positive matrices contract the hyperbolic distance in the positive cone of $\mathbb{R}^N$. Since the matrices corresponding to the maps $f_{\star n}$ are uniformly bounded in sizes and entries, this contraction is uniform. Because of this uniform contraction the set $\mathcal{M}(\Omega_T, \omega)$ is one dimensional.

Remark 5.8 *If it is easy to construct perfect tilings in dimension $d = 1$ which are not uniquely ergodic (see for instance [21]), this question remains unclear in dimension $d \geq 2$.*

Let us finish this section by giving a more specific interpretation of the Ruelle-Sullivan current associated with a transverse invariant measure in the case of tilings. Since the Ruelle-Sullivan current vanishes on exact d -differential forms, it acts on the de-Rham cohomology group $H_{DR}^d(\Omega(\mathcal{F}))$. By the same standard arguments as the one developed in Section 4.1 we have:

$$H_{DR}^d(\Omega(\mathcal{F})) = \lim_{\rightarrow} H_{DR}^d(\mathcal{F}).$$

In other words, every co-homology class $[\omega]$ in $H_{DR}^d(\Omega(\mathcal{F}))$ is the direct limit of co-homology classes $[\omega_n]$ in $H_{DR}^d(\Omega(\mathcal{F}))$. For n big enough, we have the relation:

$$\mathcal{C}_{\mu^t}([\omega]) = \sum_{i=1}^{p(n)} \frac{\mu(\pi_n^{-1}(F_i))}{\lambda_d(F_i)} \int_{F_i} \omega_n.$$

Let I_n be the standard isomorphism $I_n : H_{DR}^d(B_n) \rightarrow H^d(B_n, \mathbb{A})$ (where $\mathbb{A} = \mathbb{R}$ or $\mathbb{C}$ according to the fact that the forms have coefficients in $\mathbb{R}$ or $\mathbb{C}$) defined by:

$$\langle I_n([\omega]), c \rangle = \int_c \omega,$$

for every cycle c in $H_d(B_n, \mathbb{A})$.

As we already suggested it will be important for the proof of the gap-labelling theorem to consider the integral co-homology classes. The set of integral classes on B_n is defined by:

$$H_{DR}^{d,int}(B_n) = I_n^{-1} H^d(B_n, \mathbb{Z}),$$

it is the set of classes that take integer values on integers cycles (i.e; cycles in $H_d(B_n, \mathbb{Z})$). We define:

$$H_{DR}^{d,int}(\Omega(\mathcal{F})) = \lim_{\rightarrow} H_{DR}^{d,int}(\mathcal{F}).$$

From Corollary 4.4 and Remark 5.4 we get:

Proposition 5.9

$$\mathcal{C}_{\mu^t}(H_{DR}^{d,int}(\Omega(\mathcal{F}))) = \int_{\Omega_{T,Y}} d\mu^t \mathcal{C}(\Omega_{T,Y}, \mathbb{Z})$$

where $\mathcal{C}(\Gamma_T, \mathbb{Z})$ is the set of continuous functions on $\Omega_{T,Y}$ with integer values.

6 C^* -algebras, K -theory and Gap-Labeling

In the previous section we have seen how the description of the continuous hull Ω_T of a perfect tiling of $\mathbb{R}^d$ in terms of expanding flattening sequences has been powerful in order to describe the ergodic properties of the $\mathbb{R}^d$ -action on Ω_T . In this section we are going to use the same tools to analyze the K -theory of Ω_T .

Let us remind some very elementary facts of classical “topological” K -theory (see for instance [2]).

For any Abelian semigroup $(A, +)$ with zero 0, there is a canonical way to associate with A an Abelian group $(K(A), +)$ (also called de *Grothendieck* group of A) which satisfies a natural universal property. A simple way to construct $K(A)$ is as follows. Consider the product semigroup $A \times A$ and let Δ be its diagonal. The cosets $[(a, b)] = (a, b) + \Delta$ make a partition of $A \times A$ and we can define on the cosets set $K(A) = A \times A / \Delta$ the operation $[(a, b)] + [(c, d)] = [(a + c, b + d)]$. Note that $[(a, b)] + [(b, a)] = [(0, 0)]$, hence $K(A)$ an Abelian group.

$$\alpha : A \rightarrow K(A) \quad \alpha(a) = [a] = [(a, 0)]$$

is a semigroup homomorphism which satisfies the following universal property: for any group G and semigroup homomorphism $\gamma : A \rightarrow G$ there exists a unique homomorphism $\chi : K(A) \rightarrow G$ such that $\gamma = \chi\alpha$. If A satisfies the

cancelation rule (if $a+b = c+b$ then $a = c$), then α is injective. Let us denote by $K^+(A)$ the image of α . This is a “positive cone” in $K(A)$ (with respect to the $\mathbb{Z}$ -module structure of $K(A)$) which generates the whole $K(A)$. Every $[(a, b)]$ is of the form $[(a, b)] = [a] - [b]$. If for every $a, b \in A$, $a + b = 0$ if and only if $(a, b) = (0, 0)$, then $K^+(A) \cap -K^+(A) = \{0\}$, hence the relation $[a] - [b] \geq [a'] - [b']$ iff $[a] - [b] - [a'] + [b'] \in K^+(A)$ makes $K(A)$ an ordered group.

Let X be a *compact* topological space. Apply the above construction to the semigroup of isomorphism classes of complex vector bundles on X with the operation given by the direct sum $\oplus$. The class of the unique rank 0 vector bundle is the 0 of this semigroup. The resulting Abelian group is denoted $K^0(X)$. So each element of $K^0(X)$ is of the form $[E] - [F]$, where E and F are (classes of) vector bundles on X ; $[E] = [F]$ iff there exists G such that $E \oplus G = F \oplus G$. $K^+(X) = (\{[E] - [0]\})$ and it is the positive cone of an actual order on $K(X)$. On the other hand the semigroup does not satisfies the cancelation rule.

Since X is compact, for every F there exist $n \in \mathbb{N}$ and G such that $[F] \oplus [G] = \epsilon^n$, where ϵ^n denotes the (class of) trivial vector bundle of rank n . Hence $[E] - [F] = [E \oplus G] - [\epsilon^n]$, so that each element β of $K^0(X)$ is of the form $\beta = [H] - [\epsilon^n]$, for some $n \in \mathbb{N}$. If n_0 is the minimum of such n , then $\beta \geq 0$ iff $n_0 = 0$. Moreover if $G \oplus G' = \epsilon^m$, then $E \oplus G = F \oplus G$ implies that $E \oplus \epsilon^m = F \oplus \epsilon^m$. We can summarize this remark by saying that $[E] = [F]$ if and only if E and F are *stably equivalent*.

If $\mathcal{C}(X)$ denotes the ring of continuous complex valued functions on X , we can associate with each vector bundle E on X the $\mathcal{C}(X)$ -module of the sections of E , $\Gamma(E)$. This is in fact a functor Γ from the category $\mathcal{B}$ of vector bundles over X to the category $\mathcal{M}$ of $\mathcal{C}(X)$ -modules. The functor Γ induces an equivalence between the category $\mathcal{T}$ of trivial vector bundles to the category $\mathcal{F}$ of free $\mathcal{C}(X)$ -modules of finite rank. As X is compact, the fact that for every bundle E there exists a bundle G such that $E \oplus G$ is trivial, means that the category of vector bundles over X coincides with the sub-category $Proj(\mathcal{T})$ (which a priori is smaller) generated by the images of *projection operators* on trivial bundles. $Proj(\mathcal{F})$ is defined in a similar way. Hence Γ establishes an equivalence between $Proj(\mathcal{T})$ to $Proj(\mathcal{F})$ which is by definition the category of *finitely-generated projective* $\mathcal{C}(X)$ -modules. The construction of $K^0(X)$ can be rephrased in terms of projective modules instead of vector bundles and in such a case it is denoted $K_0(\mathcal{C}(X))$.

The Chern classes of a bundle depends only on its equivalence class β in $K^0(X)$, we denote them by $c_i(\beta) \in H^{2i}(X, \mathbb{Z})$. If $f : X \rightarrow Y$ is a continuous map between compact spaces, then there is a natural map $f_k^* : K^0(Y) \rightarrow K^0(X)$, such that for any $\beta \in K(Y)$, $c_i(f^*(\beta)) = f^*(c_i(\beta))$, where $f^* : H^{2i}(Y, \mathbb{Z}) \rightarrow H^{2i}(X, \mathbb{Z})$ is the natural map induced by f on co-homology.

The group $K^1(X)$ denotes by definition the group $K^0(SX)$, where SX is the *reduced suspension* of X , i.e. the quotient space $X \times \mathbf{S}^1 / \{x\} \times \mathbf{S}^1 \cup X \times \{p_1\}$ where x is a marked point in X and p_1 a marked point in the circle $\mathbf{S}^1$ (see [2]). Let $p : X \times \mathbf{S}^1 \rightarrow SX$ be the natural projection. If $f : X \rightarrow Y$ is as above, $(f \times id) : X \times \mathbf{S}^1 \rightarrow Y \times \mathbf{S}^1$ induces a continuous map $Sf : SX \rightarrow SY$ which can be used to define $f_k^* : K^1(Y) \rightarrow K^1(X)$.

Let us now come back to the context of tilings. Since K^0 and K^1 have a good behavior with respect to projective limits, we associate with an EFS $\mathcal{F} = \{(B_n, f_n)\}_{n \in \mathbb{N}}$, the direct sequence $K^i(\mathcal{F}) = \{(K^i(B_n), (f_n^*)_n)\}_{n \in \mathbb{N}}$, $i = 0, 1$. We have:

$$K^i(\Omega_T) = \lim_{\rightarrow} K^i(\mathcal{F}) .$$

The last statement can be understood also in terms of projection operators. Let $\psi_n : \Omega_T \mapsto B_n$ be the canonical map defined through the projective limit. Therefore any continuous function $f \in \mathcal{C}(\Omega_T)$ is the uniform limit of a sequence of functions $f_n \in \mathcal{C}(B_n)$ namely $\lim_{n \uparrow \infty} \|f - f_n \circ \psi_n\| = 0$, the norm being the sup-norm. In particular, any projection $P \in \mathcal{C}(\Omega_T) \otimes M_N(\mathbb{C})$ is equivalent to a projection in $\mathcal{C}(B_n) \otimes M_N(\mathbb{C})$ for at least one $n \in \mathbb{N}$, which means that the K groups of Ω_T is the inductive limit of the K -groups of the B_n 's.

Let us remind now some facts about the C^* -algebra $\mathcal{A} = \mathcal{A}_T$ which represents the deep non-commutative geometric structure of our dynamical system Ω_T . The C^* -algebra $\mathcal{A}$ is the crossed product C^* -algebra $\mathcal{C}(\Omega_T) \rtimes \mathbb{R}^d$. Let $\mathcal{A}_0$ be the dense sub algebra made of continuous functions on $\Omega_T \times \mathbb{R}^d$ with compact support. If μ is an $\mathbb{R}^d$ -invariant probability measure on Ω_T , there is a canonical trace $\mathcal{T}_\mu$ defined on $\mathcal{A}_0$ by $\mathcal{T}_\mu(A) = \int_{\Omega_T} d\mu(\omega) A(\omega, 0)$. The main object of the *gap-labelling* question concerns the trace $\mathcal{T}_\mu(P)$ of a projection $P \in \mathcal{A}$.

The definition of K_0 recalled above in the classical topological setting can be generalized to non commutative algebras. Recall [13] that two projections $P, Q \in \mathcal{A}$ are equivalent whenever there is an element $U \in \mathcal{A}$ such that

$P = UU^*$ and $Q = U^*U$. If P and Q are orthogonal to each other, the equivalence class of their direct sum $[P \oplus Q]$ depends only upon $[P]$ and $[Q]$, leading to the definition of the addition $[P] + [Q] = [P \oplus Q]$. To make sure that two projections can be always be made mutually orthogonal modulo equivalence, $\mathcal{A}$ must be replaced by $\mathcal{A} \otimes \mathcal{K}$. Here $\mathcal{K}$ is the C^* -algebra of compact operators on a Hilbert space with a countable basis. It can also be defined as the smallest C^* -algebra containing the increasing sequence of finite dimensional matrices: $\mathcal{K} = \varinjlim M_n(\mathbb{C})$, where the inclusion of M_n into M_{n+m} ($0 < m$) is provided by

$$A \in M_n \mapsto i_{n,m}(A) = \begin{bmatrix} A & 0 \\ 0 & 0_m \end{bmatrix} \in M_{n+m}$$

The group $K_0(\mathcal{A})$ is the group generated by formal differences $[P] - [Q]$ of equivalent classes of projections in $\mathcal{A} \otimes \mathcal{K}$. Then two equivalent projections have the same trace, and since the trace is linear, it defines a group homomorphism $\mathcal{T}_{\mu,*} : K_0(\mathcal{A}) \mapsto \mathbb{R}$ the image of which are called the *gap labels*. Together with K_0 , there is K_1 which is defined as the equivalence classes, under homotopy, of invertible elements in $\varinjlim GL_n(\mathcal{A})$. In the case of $K^1(\mathcal{C}(X))$ this is equivalent to the previous definition in terms of reduced suspensions. A standard result due to R. Bott [13] claims that $K_1(\mathcal{A})$ is isomorphic to $K_0(\mathcal{A} \otimes \mathcal{C}_0(\mathbb{R}))$ and that $K_1(\mathcal{A} \otimes \mathcal{C}_0(\mathbb{R}))$, which is then nothing but $K_2(\mathcal{A}) = K_0(\mathcal{A} \otimes \mathcal{C}_0(\mathbb{R}^2))$, is actually isomorphic to $K_0(\mathcal{A})$ (Bott's periodicity theorem). Both K_0 and K_1 are discrete Abelian groups. They are countable whenever $\mathcal{A}$ is separable. An important property of $K(\mathcal{A}) = K_0(\mathcal{A}) \oplus K_1(\mathcal{A})$ is that it defines a covariant functor which is continuous under taking inductive limits. Namely any $*$ -isomorphism $\alpha : \mathcal{A} \mapsto \mathcal{B}$ between C^* -algebras induces a group homomorphism α_* defined by $\alpha_*([P]) = [\alpha(P)]$ for K_0 and similarly for K_1 . Moreover, $K(\varinjlim \mathcal{A}_n) = \varinjlim K(\mathcal{A}_n)$.

The Thom-Connes theorem [13, 15] states that there is a group isomorphism ϕ_d between $K_0(\mathcal{C}(\Omega_T) \rtimes \mathbb{R}^d)$ and $K_d(\mathcal{C}(\Omega_T))$ (with $K_i = K_{i+2}$ by Bott's periodicity theorem). Actually the trace action makes this isomorphism more explicite:

Theorem 6.1 [14] *Let P be a projection in $\mathcal{A}$ and let $[P]$ its class in $K_0(\mathcal{A})$. (i) If d is odd, let U be a unitary element of $\mathcal{C}(\Omega_T) \otimes \mathcal{K}$ representing $\phi_d([P])$. Let η be the d -form in $H_{DR}^d(\Omega(\mathcal{F}))$:*

$$\eta = \text{Tr} \left((U^{-1}dU)^d \right) .$$

(ii) If d is even, let $Q_{\pm}$ be a pair of projections of $\mathcal{C}(\Omega_T) \otimes \mathcal{K}$ with $\phi_d([P]) = [Q_+] - [Q_-]$. Let η be the d -form in $H_{DR}^d(\Omega(\mathcal{F}), \mathbb{C})$

$$\eta = \text{Tr}((Q_+ dQ_+ \wedge dQ_+)^{d/2}) - \text{Tr}((Q_- dQ_- \wedge dQ_-)^{d/2}) .$$

Then, in both cases, and for a suitable (see later) normalization constant k_d , $k_d \eta$ is in $H_{DR}^d(\Omega(\mathcal{F}), \mathbb{Z})$ and

$$\mathcal{T}_{\mu}(P) = \mathcal{C}_{\mu^t}([k_d \eta]).$$

Combining the previous result with Proposition 5.9, we get:

$$\mathcal{T}_{\mu}(K_0(\mathcal{A})) \subset \int_{\Omega_{T,Y}} d\mu^t \mathcal{C}(\Omega_{T,Y}, \mathbb{Z}).$$

The converse inclusion being a standard result, the *gap-labelling theorem* (see theorem 1.16 stated in the introduction) is proved.

The integral classes defined above are related to Chern classes. The image, say β , of $[P]$ by the Thom-Connes isomorphism can be taken either as an element of $K^0(B_n)$, when d is even, or of $K^1(B_n)$, when d is odd, for n large enough. It turns out (this is contained in the proof of the Thom-Connes theorem) that:

1) when d is even $k_d \eta$ (which actually “lives” on B_n) represents the Chern class $c_{[d/2]}(\beta) \in H^d(B_n, \mathbb{Z})$; in fact the choice of the normalization constant k_d in theorem 6.1 ensures that this class is integral. So

$$\mathcal{T}_{\mu}(P) = \langle \mu_n | c_{[d/2]}(\beta) \rangle .$$

2) When d is odd

$$\mathcal{T}_{\mu}(P) = \langle S\mu_n | c_{[(d+1)/2]}(\beta) \rangle ,$$

where $S\mu_n$ is defined as follows. For $z\mu_n \in Z_k(B_n, \mathbb{A})$, $(\mu_n \times S^1) \in Z_{k+1}(B_n \times S^1)$ (it is understood that S^1 has the usual counterclockwise orientation). By using the natural projection $p_n : B_n \times S^1 \rightarrow SB_n$, $(\mu_n \times S^1)$ induces a $(k+1)$ (singular) cycle $S\mu_n$ on SB_n , which is called the *suspension* of the cycle μ_n .

References

- [1] Anderson J. E., Putnam I., *Topological invariants for substitution tilings and their associated C^* -algebras*, Erg. Th. Dynam. Syst., **18**, (1998), 509-537.
- [2] Atiyah M. F., *K-Theory*, Harvard University Lecture Notes (1964).
- [3] Bellissard J., *Schrödinger's operators with an almost periodic potential : an overview*, Lecture Notes in Phys., n°**153**, Springer Verlag, (1982), 356-363.
- [4] Bellissard J., *K-Theory of C^* -algebras in Solid State Physics in Statistical Mechanics and Field Theory, Mathematical Aspects*, T.C. Dorlas, M.N. Hugenholtz & M. Winnink, Lecture Notes in Physics, n°**257**, (1986), 99-156.
- [5] Bellissard J., *Gap Labelling Theorems for Schrödinger's Operators*, in *From Number Theory to Physics*, Les Houches March 89, Springer, J.M. Luck, P. Moussa & M. Waldschmidt Eds., (1993), pp.538-630.
- [6] Bellissard J., Bovier A. & Ghez J.-M., *Gap labelling Theorems for One Dimensional Discrete, Schrödinger Operators*, Rev. of Math. Phys., **4**, (1992), 1-37.
- [7] Bellissard J., Contensou E., Legrand A., *K-théorie des quasi-cristaux, image par la trace: le cas du réseau octogonal*, C. R. Acad. Sci. (Paris), **t.327**, Série I, p. 197-200, (1998).
- [8] Bellissard J., Hermmann D., Zarrouati M., *Hull of Aperiodic Solids and Gap Labelling Theorems*, in *Directions in Mathematical Quasicrystals*, CRM Monograph Series, Volume **13**, (2000), 207-259, M.B. Baake & R.V. Moody Eds., AMS Providence.
- [9] Bellissard J., Kellendonk J. & Legrand A., *Gap-Labelling for three dimensional aperiodic solids*, C. R. Acad. Sci. (Paris), **t.332**, Série I, p. 521-525, (2001).
- [10] Benameur M., Oyono H., *private communication*. J.B. thanks Benameur for this information, prior to publication.

- [11] Benedetti R., Petronio C., *Lectures on Hyperbolic Geometry*, Springer (1992).
- [12] Benedetti R., Petronio C., *Branched Standard Spines of 3-Manifolds*, LNM 1653 Springer (1997).
- [13] Blackadar B., *K-Theory for Operator Algebras*, MSRI Publications, Vol.5, Springer-Verlag, (1986).
- [14] Connes A., *Cyclic cohomology and the transverse fundamental class of a foliation*, Pitman Res. Notes in Math, **123**, Longman Harlow (1986), pp. 52-144.
- [15] Connes A., *Noncommutative Geometry*, Academic Press, San Diego, (1994).
- [16] Ghys E., *Laminations par surfaces de Riemann* in *Dynamique et géométrie complexes*, Panoramas & Synthèses **8** (1999), 49-95.
- [17] Forrest A., Hunton J., *The cohomology and K-theory of commuting homeomorphisms of the Cantor set*, Erg. Th. Dynam. Syst., **19**, (1999), 611-625.
- [18] Forrest A., Hunton J., Kellendonk J., *Cohomology of canonical projection tilings*, Submitted to Commun. Math. Phys.; *Topological invariants for projection point patterns*, to appear in Memoirs of the American Mathematical Society, XXX: math.AT/0010265. *Cohomology groups for projection point patterns*, to appear in the proceedings of the International Congress of Mathematical Physics 2000. XXX: math-ph/0010050.
- [19] van Elst A., *Gap Labelling Theorems for Schrödinger's Operators on the Square and Cubic Lattice*, Rev. of Math. Phys., **6**, (1994), 319-342.
- [20] Gähler F., Kellendonk J., *Cohomology groups for projection tilings of codimension 2*, Mat. Sci. Eng., **A 294-296**, 438-440, (2000).
- [21] Gambaudo J.-M., Martens M., *Algebraic topology for minimal Cantor sets*, preprint, Dijon (2000).
- [22] Griffiths P, Harris J., *Principles of Algebraic Geometry*, Wiley (1978).

- [23] Kaminker J., Putnam I.F., *private communication*. J.B. thanks I. Putnam for this information, prior to publication.
- [24] Kellendonk J., Putnam I.F., *Tilings, C^* -algebras and K -theory*, in *Directions in Mathematical Quasicrystals*, CRM Monograph Series, Volume **13**, (2000), 177-206, M.P. Baake & R.V. Moody Eds., AMS Providence.
- [25] Lagarias J. C. and Pleasants P. A. B., *Repetitive Delone sets and quasicrystals*, preprint, math.DS/9909033.
- [26] Milnor J.W., Stasheff J.D., *Characteristic Classes*, Princeton University Press (1974).
- [27] Moore C. C., Schochet C., *Global Analysis on Foliated Spaces*, Math. Sci. Res. Inst. Publ. **No. 9**, New York; Springer-Verlag, (1988).
- [28] Pedersen G., *C^* -algebras and their Automorphism Group*, Academic Press (1979),
- [29] Pimsner M.V., Voiculescu D., *Exact sequences for K -groups and ext-groups of certain cross-product of C^* -algebras*, J. Operator Theory, **4**, (1980), 93-118.
- [30] Pimsner M.V., *Range of traces on K_0 of reduced crossed products by free groups*, in *Operator Algebras and their Connections with Topology and Ergodic Theory*, Lecture Notes in Math., n°**1132**, Springer Verlag, (1983), 374-408.
- [31] Putnam I.F., *The C^* algebras associated with minimal homeomorphisms of the Cantor set*, Pacific Journal of Mathematics, **136**, (1989), 329-352.
- [32] Radin C., *Miles of Tiles*, Student Mathematical Library, Vol 1 (Amer. Math. Soc., Providence, 1999); see also Radin C., *The pinwheel tilings of the plane*, Annals of Math. **139**, (1994), 661-702.
- [33] Sadun L., Williams R.F., *Tiling spaces are Cantor fiber bundles*, preprint (2001), University of Texas, Austin.
- [34] Senechal M., *Crystalline Symmetries : An informal mathematical introduction*, Institute of Physics, Alan Hilger, Ltd., (1990).
- [35] Spanier E.H., *Algebraic Topology*, McGraw-Hill, New-York, (1966).

- [36] Sullivan D., *Cycles for the dynamical study of foliated manifolds and complex manifolds*, Invent. Math. **89**, (1976), 225-255.
- [37] Thurston W., *Geometry and Topology of 3-manifolds*, Princeton University LN, (1979) (<http://www.msri.org/gt3m/>).
- [38] Vershik A.M., *A Theorem on Periodical Markov Approximation in Ergodic Theory*, Ergodic Theory and Related Topics (Vitte, 1981), 195-206, Math. Res., **12**, Akademie-Verlag, Berlin, (1981).
- [39] Williams R.F., *Expanding attractors*, Publ. IHES, **43** (1974), 169-203.